

December 20, 2004

Dnr 2004/3857-52

**Recent Research on Mobile Telephony and Health Risks.  
Second annual report from SSI's Independent Expert  
Group on Electromagnetic Fields.**

***SSI's Independent Group on Electromagnetic Fields, 2004***

## **Preface**

The Swedish radiation protection agency, SSI (Statens strålskyddsinstitut) has appointed an international independent expert group (IEG) for electromagnetic fields (EMF) and health. The task is to follow and evaluate the scientific development and to give advice to the SSI. The IEG will take recent major scientific reviews as starting points and in a series of annual reports consecutively discuss and assess relevant new data and put these in the context of already available information. The result will be a gradually developing risk assessment of exposure to EMF. The group began its work in the fall of 2002 and presented its first report in December 2003. This is the second annual report.

The composition of the group for the period of 2002-2004 has been:

Prof. Anders Ahlbom, Karolinska Institutet and Stockholm Center for Public Health, Sweden (chairman);

Prof. Jukka Juutilainen, University of Kuopio, Kuopio, Finland;

Dr. Bernard Veyret, University of Bordeaux, Pessac, France;

Dr. Harri Vainio, IARC, Lyon, France (currently Occupational Health Institute, Helsinki, Finland);

Prof. Leeka Kheifets, WHO, Geneva, Switzerland (currently UCLA, Los Angeles, USA);

Prof. Eduard David, University of Witten/Herdecke, Witten, Germany;

Prof. J. Malcolm Harrington, London, UK. (Resigned in August 2004)

Ass. Prof. (Docent) Maria Feychting, Karolinska Institutet, has been appointed scientific secretary to the group.

Stockholm in December 2004

Anders Ahlbom

Chairman

## **Executive Summary**

### **Symptoms**

For RF exposure the number of studies available today is too small to allow conclusions as to whether such fields can cause various symptoms. For ELF fields the situation is different: quite a number of studies have been performed. In none of these studies have subjects been able to detect fields at levels that are comparable to those at which they claim to react. So far no study has been able to prove a link between ELF- EMF and the occurrence of symptoms. Therefore, the IEG suggests that the term “electromagnetic hypersensitivity” should not be used.

### **Cognitive functions/neurophysiology**

Recent results on RF exposure and cognitive functions have not clarified the picture. Changes in cognitive functions have been reported but, as stated in a previous review, the results are inconsistent and no single clear effect on cognitive function can be identified. The new results on ELF fields have not provided any information that would alter the conclusions of the previous reviews (the earlier positive studies generally used stronger fields).

### **EEG and sleep**

Recent investigations on humans and animals have not added clear evidence of alteration of EEG and/or sleep. In any case, the relevance of minor alterations on EEG and sleep on health is not significant.

### **Memory**

While new data have not provided evidence that memory of rodents is affected by exposure to RF fields, the data are still inconclusive in humans but possible effects do not seem to be detrimental.

### **National results from the Interphone Study**

The three first reports from the Interphone Study have been presented this year. A Swedish study suggests that long term use of mobile phones increases the risk of acoustic neuroma, at the side of the head where the phone is used. The results, however, must be interpreted with caution while other groups with sufficient numbers of long term users finalize their analyses. Both this study and a Danish study reports no effect on acoustic neuroma for short term use (The Danish study has fewer long term users). For brain tumors the Swedish study does not indicate any association with mobile phone use, for short term or long term use or for any specific tumor site.

### **Combined effects of electromagnetic fields with environmental carcinogens (CEMFEC)**

Based on the results of the animal study, cocarcinogenic effects of RF fields are not likely in this experimental model and at the exposure levels chosen. This conclusion is supported by the fact that no enhancement of in vivo genotoxic effects were found and by the fact that in

vitro studies did not show enhancement of the effects of the two chemicals or any effects characteristic to non-genotoxic carcinogenesis.

## **Exposure of children to RF and ELF fields**

Physical, chemical and therapeutic agents have the potential for affecting development, depending on the nature of the agent and the timing and magnitude of the exposure. For ELF magnetic fields there are indications that children might be more sensitive; however, we lack the understanding of how, or even if, these fields might be involved in leukemogenesis. Widespread exposure to these fields is recent and very little is known about the potential sensitivity of children to RF fields. Given the paucity of data indicating a particular vulnerability of children to EMF, it may be tempting to conclude that children are not more susceptible than adults to RF exposure. However, the absence of an observed effect does not necessarily mean that exposure is harmless, especially if crucial studies focusing on children are yet to be done. Given scientific uncertainty SSI has adopted precautionary approaches for both ELF and RF which we endorse.

## ***Introduction***

The 2003 annual report of the IEG focused on RF and in particular epidemiological studies on cancer, as well as on studies on experimental carcinogenicity, heat shock proteins, and blood brain barrier effects [Swedish Radiation Protection Authority 2003]. The IEG noted in all these sections that there were various open questions but also that current research was addressing these issues attempting to confirm or replicate previous results or by looking at complementary aspects. The plan for the IEG is to revisit each of these issues when new data are available and to update the discussion and conclusions. This year we report on some new epidemiologic data, but for the other issues too little new data have emerged and further discussions of these topics will have to wait.

This annual report brings up a set of different issues. We first review the literature on a widely discussed topic, namely whether exposure to fields increases the frequency of asthenic symptoms either in a susceptible group of the population or in the general population; this is often referred to as electrical hypersensitivity. Somewhat linked to this is the possibility of cognitive effects and neurophysiological effects from EMF exposure and we review this as well as studies on EEG and sleep and memory in the present report. As in 2003, the starting point is the Stewart report from 2000 [IEGMP 2000] and the update of the Stewart report made in 2003 by NRPB's independent Advisory Group on Non-Ionising Radiation (AGNIR) [NRPB 2003]. Several European research programs have come to, or are about to come to, an end and we have included a summary and discussion of the main results of some of these programs. It is in this context we discuss some recent epidemiological data that have come out of the Interphone Study. The extent to which children might be extra sensitive to electromagnetic fields was touched upon in the previous year's report but was postponed till this year primarily to include the discussion of the WHO-conference on this topic which was held during the summer of 2004. In this context we also discuss exposure policy for children. A starting point for this discussion is IARC's classification of ELF-EMF in Group 2B "possible carcinogen to humans".

## **Symptoms**

### **Background**

The term "electromagnetic hypersensitivity" (EHS) is often used to denote a phenomenon where individuals experience adverse health effects while using or being in the vicinity of electric, magnetic, or electromagnetic field sources and devices, and when the individuals themselves attribute their symptoms to these sources and devices. There are no diagnostic criteria available, and symptoms experienced vary substantially between different individuals, but are generally non-specific with no objective signs present. The severity of the condition varies; the majority of cases present mild symptoms, but some cases experience severe problems with major consequences for work and everyday life. The scientific literature on this area has been reviewed previously by Bergqvist et al. for the European Commission [Bergqvist, et al. 1997], updated in 2000 [Bergqvist, et al. 2000], and by Levallois [Levallois 2002]. This report briefly summarizes the findings of the previous reviews, and evaluates additional studies available in the scientific literature.

It is difficult or even impossible to correctly estimate the prevalence of EHS, simply because there are no established diagnostic criteria, and the definition of the disease is largely based on each individual's own beliefs and attribution of various symptoms to different sources of electromagnetic field exposure. Therefore, assessments of the prevalence will entirely depend on the methods used to identify cases, and the types of questions asked in each specific survey. The hitherto reported prevalence of EHS varies considerably throughout the world and between reports; at the time of the review for the European Commission [Bergqvist, et al. 1997] it was most common in the Nordic countries and Germany with less than 1%, but very rare or non-existent in other countries, e.g. the UK and the Netherlands. A cross-sectional survey performed in 1997 on a random sample of the population in Stockholm, Sweden, reported a prevalence of 1.5% [Hillert, et al. 2002], and a survey made on a sample of the general population in California estimated the prevalence to 3.2% [Levallois, et al. 2002]. For reasons discussed above the validity of the reported prevalence in different studies can be questioned, and the prevalence in different countries cannot easily be compared.

Röösli et al. [Roösli, et al. 2004] report results from a Swiss descriptive study of persons complaining about symptoms that they attributed to electromagnetic fields. The most common symptoms were sleep disorders, followed by headaches, nervousness or distress, fatigue and concentration difficulties. The most common sources to which the subjects attributed their symptoms were mobile phone base stations (74%), mobile phones (36%), cordless phones (29%) and power lines (27%).

Surprisingly few etiologic studies have been made on this health problem, and the majority of them have focused on skin symptoms related to video display terminal (VDT) use. The unspecific nature of the disease is a major difficulty in this research, especially because the patient's attribution of the symptoms to sources of electromagnetic field exposure is an important part of the disease definition.

Studies on "electromagnetic hypersensitivity" can for natural reasons only be made on humans, and the available studies are either epidemiological (observational) or experimental (provocation studies). Most studies have focused on extremely low frequency (ELF) fields or VDT use, but there are also some studies on radiofrequency (RF) exposure from mobile phones or base stations.

## RF exposure and symptoms

### Epidemiological studies

To date all of the existing epidemiological studies on RF exposure and symptoms are cross-sectional, which makes them of limited value in an evaluation of whether low level RF exposure can cause various symptoms. In a cross-sectional study the exposure and the outcome are assessed simultaneously, without consideration of the time sequence of the events. In addition, all of the available studies on RF exposure ask the subjects themselves to assess both their exposure (e.g. distance to nearest base station or amount of mobile phone use) and the outcome (various symptoms), which lead to a considerable risk that the exposure assessment is influenced by the disease, or that only symptoms that an individual attribute to the RF exposure are reported. Another limitation is that the participation rates in most of the studies are low or not possible to assess, and there is a large potential for selection bias; people who experience symptoms that they attribute to mobile phones or base stations may be more prone to participate in a study investigating this particular question, than people with no such beliefs. Also the Stewart report [IEGMP 2000] acknowledged the limitations inherited in studies using a cross-sectional design.

At present there are only two studies on base station exposure and symptoms published; one from France [Santini, et al. 2002a] and one from Spain [Navarro, et al. 2003]. None of the studies have reported how subjects were selected for participation in the studies, and participation rates cannot be estimated. Participants have answered questions about various symptoms such as headaches, concentration difficulties, memory loss, fatigue, sleeping problems etc. They were also asked to estimate the distance to the nearest base station, with no independent validation. Not only is self reported distance to base stations a questionable exposure assessment method; it has also been shown that distance is a poor surrogate for RF exposure from base stations [Schuz and Mann 2000]. The Spanish study also made measurements of the exposure in the homes, but did neither report how subjects were selected for measurements nor the proportion of subjects agreeing to have measurements taken in their homes. For unknown reasons they have also excluded all participants living between 150 and 250 meters from a base station, which makes correlation coefficients between distance and exposure of little value. Both the French and the Spanish study report an increased prevalence of symptoms close to base stations, but the design limitations make it impossible to assess whether these findings are a results of bias or real effects.

Three studies have investigated the prevalence of symptoms among mobile phone users [Chia, et al. 2000; Oftedal, et al. 2000; Santini, et al. 2002b], with participation rates varying from 45% to 65%. The study by Oftedal et al. was performed to assess differences between users of analogue and digital phones. Various symptoms were reported among heavy users of mobile phones primarily among participants from Norway and to a lesser degree among participants from Sweden, but did not differ between users of different types of phones. The symptoms reported were a feeling of warmth on, around or behind the ear, headache, dizziness, fatigue, concentration difficulties. The study by Chia et al. reported a somewhat increased prevalence of headaches among mobile phone users, and a lower prevalence of concentration difficulties. None of the other types of symptoms differed between mobile phone users and others. Santini et al. found no differences in reported symptoms related to mobile phone use. The results vary considerably between studies, and it is possible that the way the study was introduced to participants, how questions were phrased, recent reports in the local media, or differences in participation rates may have influenced the results. These findings cannot be used to draw conclusions about the effect of mobile phone use on the occurrence of various symptoms.

In conclusion, the available epidemiological studies provide little information regarding the question if RF exposure from base stations or mobile phones can cause various symptoms.

### Experimental studies

Another approach to test the hypothesis that RF exposure from mobile phones or base stations can cause various symptoms is to perform an experimental study where the exposure circumstances are controlled by the investigator in a so called provocation study. In this type of study the occurrence of symptoms during periods of exposure are compared to the occurrence of symptoms during unexposed periods. The ideal design is a double blind study where neither the investigator nor the subject is aware of when the exposure is on or off, which minimizes the possibility that knowledge about the exposure will influence the assessment of the studied outcome. Many electrically hypersensitive persons claim that their symptoms occur immediately after having been close to for example an active mobile phone, and for this group of EHS persons an experimental study is probably the most valid study design. So far only three provocation studies have been performed on RF exposure and symptoms.

A Finnish research group reported results from two provocation experiments each including 48 healthy subjects [Koivisto, et al. 2001]. In one experiment, subjects were studied during two 60 min sessions 24 hours apart, where half of the subjects were exposed to GSM 900 MHz during the first session and half during the second, and the other study had two 30 min sessions following directly after each other. Subjects were not aware of when the exposure was on or off. No association between the exposure and occurrence of symptoms was found, and generally few symptoms were reported. If only a small proportion of the population are affected by exposure to mobile phones this would not have been detected in this small study of healthy volunteers.

Another Finnish study [Hietanen, et al. 2002] included 20 subjects that were selected because they claimed to be sensitive to exposure from mobile phones. The purpose was to study symptoms occurring in sensitive persons after exposure to RF fields from mobile phones, and to test their ability to detect whether the fields were on or off. The study used a double blind design and reported results from four 30 min sessions 60 min apart: one session with exposure to a 900 MHz analogue mobile phone, one with a 900 MHz digital phone, one with an 1800 MHz digital phone and one with no exposure. Study subjects were asked to report experienced symptoms immediately, and all symptoms that occurred during a session disappeared during the break. The results showed that more symptoms were reported during the sessions without exposure than when the exposure was on. The subjects were unable to detect the presence of the RF field better than chance. A limitation in the study is that the unexposed session always was one of the first two sessions; there are indications that the participants experienced more stress in the beginning of the experiment which may have hampered the ability to discover an association should one exist.

These are the only two available provocation studies of mobile phone use, and both have limitations that prevent conclusions.

A Dutch study of exposures similar to that from mobile phone base stations, the so called TNO-study, included one group of 36 EHS subjects claiming to experience symptoms in connection to GSM mobile telephony and one group with 36 healthy subjects [Zwamborn, et

al. 2003]. The groups differed in terms of age and gender distribution and therefore no comparisons could be made between the groups, only within groups for periods with and without exposure. The subjects were exposed to a 1 V/m field at 900 MHz and 1800 MHz (GSM signal), and 2100 MHz (UMTS signal). Each subject participated in three sessions one of which was unexposed, using a double blind design. Each session took 45 min including exposure, questionnaire, and break. The questionnaire estimated degree of well being from 23 assessed symptoms. Both the EHS group and the healthy volunteers reported a somewhat lower degree of well being during UMTS exposure compared to no exposure, whereas no effects were seen for periods of GSM 900 or 1800 exposures in either of the groups. This is the only available study of base station exposures and the results need to be interpreted cautiously, which is also acknowledged by the authors. The results do not explain existing reports of perceived symptoms from base stations since these symptoms are related to the second generation of mobile telephony. Additional studies are needed before any conclusions can be drawn. One such replication study is ongoing in Switzerland; the study protocol is available on the web: [www.mobile-research.ethz.ch](http://www.mobile-research.ethz.ch).

## **ELF exposure and symptoms**

Detailed reviews of these studies can be found elsewhere [Bergqvist, et al. 2000; Bergqvist, et al. 1997; Levallois 2002]. These reviews concluded that some of the epidemiological studies of skin disorders and VDT use report an excess prevalence of subjectively reported symptoms among VDT users, whereas no consistent association was seen between VDT use and objective signs. All of these studies are cross-sectional, and can therefore not be used to assess whether electromagnetic fields from the VDT can cause skin disorders. A study published after these reviews found no differences in magnetic field levels assessed over 24 h between EHS patients and healthy controls [Sandstrom, et al. 2003], but did also use a cross sectional design. The purpose of the study was to describe differences between EHS patients and healthy controls in terms of ECG, heart rate, and heart rate variability, as an indication of a dysbalance in the autonomic nervous system among EHS patients, but with no attempt to determine the cause of any differences.

Several experimental provocation studies have been performed where EHS subjects were exposed in the laboratory to test their ability to detect the electromagnetic fields. The studies used a double-blind crossover design with different exposure settings; one study exposed subjects to either a personal computer (PC) with low fields, or one with high fields ([Swanbeck and Bleeker 1989], one used a PC that was either on or off [Andersson, et al. 1996], one used a filter that reduced the fields from the VDT, a passive filter, or no filter [Oftedal, et al. 1999], and one combined exposure from a VDT with other stressors [Lonne-Rahm, et al. 2000]. Neither in studies of subjects having primarily skin symptoms in connection to VDU work, nor in studies of subjects with a more general electromagnetic hypersensitivity were EHS patients able to detect the fields. Adding additional stressors did not change this picture. Some of these studies exposed subjects in an open session to establish that they did react to the fields, but they could not detect the fields in subsequent double blind experimental sessions. In some studies, subjects reported symptoms when they believed that the fields were on, but unrelated to the actual exposure. The study of filters for reduction of the fields from the VDT found a significant decrease of symptoms during periods with filters compared to periods with no filter, regardless of whether the filter was active or passive [Oftedal, et al. 1999].

The experimental studies have been criticized for the unnatural experimental situation which may have influenced the subjects' ability to detect the fields. To take this into consideration a Swedish study was performed in the homes or workplaces of EHS subjects [Flodin, et al. 2000], with several days between sessions and a 24 h of follow-up to detect late appearing symptoms. A control group with healthy volunteers was also included. EHS subjects could not better than controls detect exposure to electric and magnetic fields.

## **Overall conclusions**

For ELF fields, quite a number of studies have been performed, and in none of these studies have EHS patients been able to detect electric or magnetic fields at levels that are comparable to those at which they claim to react.

For RF exposure the number of studies available today is too small to allow any conclusions as to whether these fields can cause various symptoms. Several studies are ongoing, and will hopefully report results within the near future.

So far no study has been able to prove a link between electromagnetic fields and the occurrence of symptoms.

Recently a workshop was organized by WHO to review, evaluate, and discuss the scientific evidence related to the question of whether electromagnetic field exposure can cause various symptoms (WHO workshop, Prague, October 2004). At this workshop, no new data was presented that would change the picture described above. It was suggested that the term electromagnetic hypersensitivity should not be used, because of the lack of evidence that EMF exposure plays a role in the occurrence of the symptoms. Instead the term idiopathic environmental intolerance (IEI) was suggested. However, the IEG does not believe that this term is any better. In fact, any term that combines exposure and health will make scientific investigations of etiology more complex. Three working groups were formed at the WHO workshop; one that will work on the definition of the outcome and intervention strategies, one that will discuss communication, and one that will address the need for future studies and research recommendations.

An epidemiological study properly designed to test the hypothesis that electric or magnetic fields can cause various symptoms in a small group of sensitive persons would need to be a very large prospective cohort study with repeated assessment of both the outcome and the exposure, which would be a huge undertaking. As long as experimental studies have not been able to establish that electromagnetic fields can trigger symptoms in EHS patients, such a study cannot be recommended.

## **Cognitive functions/neurophysiology, RF, ELF**

### **RF fields**

*"The body of evidence regarding direct RF exposure effects on cognitive function remains inconsistent. Some well-conducted studies have reported significant changes in cognitive functions due to RF exposure, while others have found no significant effects. Among those studies reporting changes, both facilitatory and inhibitory effects of RF exposure have been reported. Overall, no single clear effect of RF exposure on cognitive function can be identified"* After the above conclusions by AGNIR (2003), a few studies have been published, and are reviewed here.

Maier et al. [Maier, et al. 2004] exposed 11 volunteers to a GSM-type RF field from a programmable cellular phone held 4 cm from the left ear (unknown whether 900 or 1800 MHz). An auditory discrimination task was applied and participant's current 'Order Threshold' (OT) value was determined. The OT was defined as the minimum time needed to a) recognize that two auditory stimuli are presented just separately and b) to decide without error on what side the first of both stimuli was presented. Following a first test cycle, the volunteers had to relax for 50 min while being, or not being, exposed to pulsed electromagnetic fields. Subsequently, the test was repeated. Neither the participants nor the experimenter were aware of when the exposure was on or off. Data acquired before and after the resting phase were compared from both experimental conditions. Nine of the 11 test participants showed worse results in their auditory discrimination performance upon field exposure as compared with control conditions ( $p=0.0105$ ).

Curcio et al. [Curcio, et al. 2004] investigated the time-course of electromagnetic field (EMF)-induced effects on human cognitive and behavioral performance and on tympanic temperature. Subjects were randomly assigned to two groups, exposed to a GSM-900 signal before the testing session, or to the same signal during the data collecting session (maximum local SAR of 0.5 W/kg). Using a double-blind paradigm, subjects were tested on four performance tasks: an acoustic simple reaction time task, a visual search task, an arithmetic descending subtraction task and an acoustic choice reaction time task. Moreover, tympanic temperature was collected five times during each session. Results indicated an improvement of both simple and choice reaction times and an increase of local temperature on the exposed region under the active exposure. No effects were seen on the visual search task or the arithmetic descending subtraction task. There was a clear time-course of the reaction time and temperature data, indicating that performance and physiological measures need a minimum of 25 min of EMF exposure to show appreciable changes.

The so called TNO study from the Netherlands described in the previous chapter on symptoms also examined the effects of RF signals from mobile phone base stations on cognitive functions [Zwamborn, et al. 2003]. Cognitive functions were measured during exposure. In the cognitive function tests (reaction time, memory comparison, dual-tasking, visual selective and filtering irrelevant information), statistically significant differences were seen more often than should occur by chance (due to statistical noise), but there was no consistent pattern of results across the three signals, the different cognitive tasks or the two study groups. Independent replication of these findings would be useful.

In summary, recent results on RF exposure and cognitive functions have not clarified the picture. Changes in cognitive functions have been reported but, as stated by AGNIR (2003), the results are inconsistent and no single clear effect on cognitive function can be identified.

## **ELF fields**

Possible effects of low frequency fields on human cognition and performance were reviewed in the recent ICNIRP (2003) report, and concluded as follows: *“In summary, some changes of magnetic fields on reaction time and accuracy have been reported but the effects were not consistent between studies. For memory and vigilance, there are essentially no reported effects. Further studies are required to clarify these issues [possible effects on reaction time and accuracy]. In addition, the nature of these effects makes it difficult at present to assign them any importance with regard to health impairment”*. Similar conclusions were reached in a recent review [Crasson 2003]: *“The psychophysiological and cognitive studies indicate that we cannot exclude the possibility of 50-60 Hz weak magnetic field exposure on human cognitive processes. The observed effects, however, are small...Effects are inconsistent, subtle, transitory and specific to some aspects of cognitive functioning, without a clear dose-response relationship, and difficult to reproduce.”*

After the completion of the 2003 reviews by ICNIRP and Crasson, Kurokawa et al. [Kurokawa, et al. 2003] examined acute effects of 50-Hz magnetic fields on cognitive performance in humans. Twenty young subjects were exposed to circularly polarized 20- $\mu$ T fields for 55 min, during which they performed four cognitive performance tests for evaluation of simple reaction time, time and accuracy of choice reaction, time perception, and figure perception. None of the subjects detected the existence of the field, and no effects on performance in the tests were observed.

The new results on ELF fields have not provided any such information that would alter the conclusions of the previous reviews (the earlier positive studies generally used stronger fields than Kurokawa et al.).

## **EEG and sleep**

EEG and sleep studies are here considered together, as EEG monitoring is being used as a major tool in sleep research and several of the results are related to EEG during sleep. A few research and review papers mainly dealing with RF exposure have recently been published that are reviewed below:

In Switzerland, Huber et al. had shown effects on waking regional cerebral blood flow (rCBF) and on waking and sleep electroencephalogram (EEG) in humans [Huber, et al. 2002]. Exposure to GSM-900 was on one side of the head during 30 min and the spatial peak SAR in the head was 1 W/kg. The authors used PET<sup>1</sup> to monitor rCBF following the exposure. They also recorded the quality of sleep during night-time sleep. The rCBF increased on the side of

---

<sup>1</sup> positon emission tomography

the head exposed and the EEG power was altered. However, continuous signals did not elicit this effect.

More recently, the same group exposed the volunteers either during sleep or during the waking period preceding sleep [Huber, et al. 2003]. In the first experiment, subjects were exposed intermittently during an 8-hour night-time sleep period and, in the second experiment, on one side of the head for 30 min before a 3-hour daytime sleep period. Compared to the control condition with sham exposure, the spectral power of the non-REM<sup>2</sup> sleep EEG was initially increased in the 9-14 Hz range in both experiments. Unilateral exposure during waking induced a similar effect in both hemispheres of the brain. Some characteristics of the sleep pattern were altered. The authors interpret the effects as originating from a structure below the cortex such as the thalamus which was similarly exposed at around 0.1 W/kg.

These two studies have been carefully planned and performed and have yielded some interesting data. The amplitude of the effects remains small and of unknown consequences for health. However, further investigations will help ascertain the existence of the effects and their possible mechanism.

In Finland, Krause and co-workers recently published the results of a replication of their own work [Krause, et al. 2004]. The effects of GSM-900 exposure on the event-related desynchronization/synchronization (ERD/ERS) of some of the EEG frequency bands were studied in 24 normal subjects performing an auditory memory task. In this double-blind study, all subjects performed the memory task both with and without exposure in a counterbalanced order. The authors were not able to replicate the findings from their earlier study. Also, the effect of exposure on the number of incorrect answers in the memory task was inconsistent. They concluded that GSM-900 effects on the EEG and on the performance of memory tasks is variable and not easily replicable for unknown reasons.

In the three human studies described below, various approaches were used for exposure and EEG monitoring but several shortcomings, in particular in terms of dosimetry, make it difficult to agree with the authors' conclusions about the existence and relevance of effects of exposure on EEG:

- In the Australian study of D'Costa and co-workers, EEG recordings from 10 awake subjects were taken during exposure to a GSM phone positioned behind the head, the antenna pointing toward the head [no estimation of SAR was given for this unusual exposure situation] [D'Costa, et al. 2003]. Two experimental trials were conducted. In the first trial, the GSM mobile phone had its speaker disabled and configured to transmit at full-radiated power. In the second trial, the mobile phone was in active standby mode. For each trial, subjects were exposed under single-blind conditions in 5-min intervals to a randomized, interrupted sequence of 5 active and 5 sham exposures. The average EEG band power in active exposure recordings was compared to corresponding sham recordings. The EEG alpha (8-13 Hz) and beta (13-32 Hz) bands showed differences when the full-power mode was used.
- Kramarenko and Tan, in Ukraine, recorded EEG changes during exposure of volunteers to a GSM phone on stand-by [Kramarenko and Tan 2003]. They claim to have suppressed the interference caused by emission from the phone and observed changes in EEG patterns: after 20-40 s, a slow-wave delta activity (2.5-6.0 Hz) appeared in areas on the side of the phone. After turning off the mobile phone, slow-wave activity progressively disappeared. They observed similar changes in children, but the slow-

---

<sup>2</sup> REM: rapid eye movement i.e. "dream phase" of sleep

waves with higher amplitude appeared earlier in children. According to the authors, these results suggested that cellular phones may induce abnormal slow waves in EEG of awake persons. However, it is most surprising that they were able to record slow-waves in awake volunteers and to transmit the signal by telemetry. Moreover, the dosimetry was not well described. Overall, this study does not bring new evidence on EEG effects of exposure.

- In the Estonian study of Hinrikus and co-workers, 20 healthy volunteers were exposed to 450-MHz microwaves with 7-Hz on-off modulation [Hinrikus, et al. 2004]. The field power density at the scalp was very low ( $0.16 \text{ mW/cm}^2$ , at the scalp 10 cm away from the antenna, i.e.  $9.5 \text{ mW/kg}$  in the brain). Microwave stimulation caused changes in EEG in the frontal region. Changes varied strongly from subject to subject but overall, alterations of EEG were not statistically significant.

Few results have been reported on animals in recent years. Marino and co-workers studied the potential nonlinear changes in brain electrical activity due to cell phone exposure in rabbits [Marino, et al. 2003]. When the animals were exposed to the radiation from a mobile telephone (800 MHz, 0.6 W maximum emitted power), under conditions that simulated normal human use, the EEG was affected in nine of the ten animals studied. The effect occurred 100 ms after the onset of exposure and lasted approximately 300 ms. Absorption of the radiation by the EEG electrodes could not account for the observed effect. No effect was seen when exposure of the brain was minimized by moving the antenna from the head to the chest. The authors concluded that exposure to a standard mobile telephone can alter brain function. However, great caution again must be taken to interpret these data obtained during exposure on a short time scale. Extrapolation to human health is questionable even if the effect were real.

Three reviews have been published recently:

- Cook and co-workers dealt mainly with ELF exposure (and ELF-modulated RF) [Cook, et al. 2002]. The conclusion for ELF was that “The investigation of weak (below 500  $\mu\text{T}$ ), extremely low frequency (ELF, 0-300 Hz) magnetic field (MF) exposure upon human cognition and electrophysiology has yielded incomplete and contradictory evidence that MFs interact with human biology.”
- Hamblin and Wood reviewed 14 papers on EEG and sleep related to GSM exposure [Hamblin and Wood 2002]. They concluded that, in general, outcomes have been inconsistent and comparison between individual studies was difficult. However, enhanced EEG alpha-band power has been noted in several of the studies, a phenomenon also observed in some animal studies. Since this alteration had been found in some ELF work, the authors suggested that the effect could be caused by exposure to the ELF magnetic field produced by the phone battery. Effects have been reported to occur both during exposure and up to 1 hour or so after cessation of exposure. It is however difficult to agree with the authors about the possible role of the magnetic field produced by the phone in eliciting the effects as there are no well-established effects of ELF magnetic fields on EEG (see above) and as the magnetic field is produced only very close to the phone, which in some of the investigations on EEG was not even in contact with the head.
- The review by D’Andrea and co-workers is part of an ICES series of “white papers” on microwaves and health [D’Andrea, et al. 2003]. The authors concluded that effects reported on sleep EEG are more likely to involve non REM alpha waves, compared to other bands. Serious deficiencies in EEG human studies using EMF exposure included poor dosimetry making it difficult to compare results among experiments and

laboratories. The authors concluded that *“no conclusions can be drawn from the present EMF–EEG research. If EMF–EEG research continues, the simplest of paradigms, and traceable dosimetry must be considered. The relation of the antenna to the brain must be defined in stereotaxic coordinates and the SAR fully mapped in the brain experimentally and numerically.”*

## Conclusion on EEG and sleep

Recent investigations on humans and animals have not added clear evidence of alteration of EEG and/or sleep. In any case, the relevance of minor alterations on EEG and sleep on health is not significant. Further investigations on sleep are ongoing and may bring some new information on these subtle effects.

## Memory

From the AGNIR report [NRPB 2003]: *“The evidence from several laboratories indicates that changes may be induced in cholinergic activity in the brain following intense exposure. Such changes might predict effects on spatial learning and memory, but the evidence for this is equivocal: two studies from one laboratory have reported deficits in performance of spatial memory tasks using pulsed microwaves. But field-dependent changes were not confirmed in two independent studies using GSM signals. In addition, significant deficits on the performance of a related task were seen only when exposure increased body temperature by 2°C. Nevertheless, few tasks and exposure conditions have been examined and the available results do not rule out the possibility that microwaves may engender subtle cognitive or behavioural changes in immature or adult animals.”*

## Animal studies

Beyond the few studies that have concerned human memory, most of the research activity has been devoted to the replication and conformation of the results of the Lai group obtained on rats exposed whole-body to pulsed RF fields at 2.45 GHz [Lai, et al. 1994; Wang and Lai 2000]. See also the next chapter on results from some recent studies within the European Union 5<sup>th</sup> research program.

Both the groups of Cobb and Cassel who did replication studies using the exposure system of Lai found no evidence of impairment of memory in rats exposed at 2.45 GHz [Cassel, et al. 2004; Cobb, et al. 2004]. In parallel, the groups of Edeline and Sienkiewicz found no effect on spatial memory in rodents exposed using GSM signals [Dubreuil, et al. 2002; Dubreuil, et al. 2003; Sienkiewicz, et al. 2000].

The conclusion is thus today that there is no evidence of an effect of exposure to RF on memory of rodents. Since it was not clear what was the behavioural parameter tested in the Lai experiments (spatial memory, visual memory, performance, anxiety, etc.), further studies have recently been performed by the Cassel group on anxiety using an elevated plus maze and no effects were seen [Cosquer, et al. 2004].

However, Lai has recently published data on the effect of magnetic field noise that suppresses the effect of microwaves on rodent memory [Lai 2004]. It is thus surprising that the effect seems to be present in his laboratory and not in others and even more that “magnetic noise”

suppresses the effect, as there is no solid evidence today that such ELF fields can remove bioeffects of ELF fields.

### Human studies

Edelstyn and Oldershaw found that human volunteers exposed to mobile phones showed improvement in the performance of three cognitive function tests: (i) immediate verbal memory capacity, (ii) immediate visual spatial working memory capacity, and (iii) sustained attention [Edelstyn and Oldershaw 2002].

Two recent RF human cognitive studies have been performed [Haarala, et al. 2003; Krause, et al. 2004]; both were double-blind studies that failed to replicate some earlier findings [Koivisto, et al. 2000].

In conclusion, while there is no evidence that memory of rodents is affected by exposure to RF fields, the data are still inconclusive in humans, even if effects are of small amplitude and do not seem to be detrimental.

## **Results from some recent studies within the European Union 5<sup>th</sup> research program**

### **National results from the INTERPHONE study**

In the beginning of this year results were reported from a Danish case-control study of acoustic neuroma and mobile phone use [Christensen, et al. 2004], that is part of the INTERPHONE study. Information about mobile phone use was collected through personal interviews with 106 cases and 112 randomly selected controls matched on age and sex. Persons who were regular mobile phone users did not have an increased risk of acoustic neuroma (relative risk=0.9; 95% CI 0.5-1.6). The relative risk did not increase regardless of duration of use, amount of use, or time since first regular use. The number of long-term users was, however, small; only two acoustic neuroma cases and 15 controls had started to use a mobile phone more than ten years prior to diagnosis (RR =0.2; 95% CI 0.04-1.1).

During the fall, results from the Swedish part of the INTERPHONE study were reported [Lönn 2004; Lönn, et al. 2004]. The analyses were based on 644 brain tumor cases, 148 acoustic neuroma cases, and 674 randomly selected controls stratified on age, sex, and geographic region (604 controls in the acoustic neuroma analyses). Mobile phone use that started less than 10 years prior to diagnosis was not associated with an increased risk of any of the studied tumor types, whereas mobile phone use that started at least 10 years prior to diagnosis was associated with an increased risk of acoustic neuroma (RR=1.9; 95% CI 0.9-4.1, based on 14 exposed cases), which was confined to the side of the head where the phone was usually held (RR=3.9; 95% CI 1.6-9.5). For brain tumors, no indications of increased risks were found, regardless of duration of mobile phone use, or if analyses were restricted to the tumor location where the exposure is highest.

The results for short term mobile phone use in these two studies are consistent with the majority of previous reports [Auvinen, et al. 2002; Dreyer, et al. 1999; Inskip, et al. 2001; Johansen, et al. 2001; Muscat, et al. 2002; Muscat, et al. 2000], although the numbers of long term users in previous studies have been too small. The only previous studies that have consistently observed increased risks found risk elevations after a short duration of mobile phone use, and for several different types of tumors [Hardell, et al. 2002; Hardell, et al. 2001], which is not consistent with the studies described above.

### **Combined Effects of Electromagnetic Fields with Environmental Carcinogens (CEMFEC)**

In the CEMFEC project, coordinated by the University of Kuopio and funded by the 5<sup>th</sup> Framework Research Program of the European Union, cocarcinogenic effects of low-level GSM-type radiofrequency (900 MHz) radiation were evaluated in a 2-year animal study conducted in compliance with the OECD Principles of Good Laboratory Practice (GLP). The known carcinogen used to induce cancer was the drinking water mutagen (3-chloro-4-(dichloromethyl)-5-hydroxy-2(5H)-furanone, MX), which is known to act as a multi-site carcinogen in Wistar rats. Female Wistar rats (72 animals/group) were exposed to RF fields at two exposure levels (Specific Absorption Rates of 0.3 W/kg and 0.9 W/kg) for 2 h per day on

5 days per week. At the end of the 2-year study, tissue samples were collected for histopathology.

After 3, 6 months and 2 years of exposure, blood samples were collected (20 animals/group) for genotoxicological assays. Brain and liver tissue samples were taken for the same purpose at the end of the study. The comet assay was used to detect DNA damage and the micronucleus test to detect chromosomal aberrations (structural chromosome damage) and genome mutations (aneuploidy induction).

Also *in vitro* studies using two cell lines were conducted to investigate whether RF electromagnetic fields affect selected cellular events that have been proposed as mechanisms for non-genotoxic carcinogenesis or cocarcinogenesis. In addition to MX, another environmental chemical (the fungicide Vinclozolin, or 3-(3,5-dichlorophenyl)-5-methyl-5-vinylloxazolidine-2,4-dione) was tested for combined effects with RF radiation. The mammalian cell lines NIH3T3 and L929 were used. Several endpoints were evaluated, including oxidative stress, cell proliferation, cell cycle analysis, apoptosis, mitochondrial membrane potential modifications and gene expression (proto-oncogens: c-fos, c-jun and c-myc). The SAR levels were similar to those of the animal study (0.3 W/kg and 1.0 W/kg).

The results of the animal study confirmed the carcinogenic effect of MX, but the RF exposures did not affect statistically significantly the incidence of any tumour type.

RF exposure did not induce genotoxic effects in blood, liver or brain. There were indications of a slightly decreased genotoxic response in the RF-exposed animals in blood cells collected at 3 and 6 months. These differences lost statistical significance after 24 months of exposure. RF exposure did not affect the level of DNA damage in liver or brain, either.

The *in vitro* experiments did not show any consistent effects on oxidative stress, apoptosis, mitochondrial membrane potentials or expression of selected oncogenes. However, both RF alone and in combination with the chemical exposures suppressed cell growth. This effect was dependent on exposure time (seen only after 24 h of exposure).

Based on the results of the animal study, cocarcinogenic effects of RF fields are not likely in this experimental model and at the exposure levels chosen. This conclusion is supported by the fact that no enhancement of *in vivo* genotoxic effects were found, and by the fact that *in vitro* studies did not show enhancement of the effects of the two chemicals or any effects characteristic to non-genotoxic carcinogenesis. In fact, according to current understanding of carcinogenesis, both the suggestive “protective” effect of the RF exposures against DNA damage and the slightly decreased proliferation in RF-exposed cells (if confirmed in further studies) should reduce rather than increase cancer risk.

The fact that SAR values higher than 1 W/kg were not used may be considered a limitation of the project. However, use of higher values might have caused thermal effects and thus problems with the interpretation of the results.

In summary, the results of the CEMFEC study should be interpreted together with other completed and ongoing studies using different experimental models and varying exposure levels.

## **Perform B: *In-vitro* and *in-vivo* Replication Studies Related to Mobile Telephones and Base Stations**

The Perform B research programme lasted two and a half years and just ended. It was entitled “*In-vitro* and *in-vivo* Replication Studies Related to Mobile Telephones and Base Stations.” The PIOM laboratory in Bordeaux, France was the co-ordinator of the programme. Its objective was to provide key research results which, when combined with current and past studies on the effects of radiofrequency radiation (RFR) on living systems, would contribute to an adequate database for health risk evaluation by public health authorities. It addressed specifically *in vitro* and *in vivo* replication studies recommended by the WHO agenda of 1999.

Two different biology laboratories were involved in each of the three proposed studies. The main emphasis was thus on replication rather than on mechanistic investigations. Several exposure systems were designed or adapted: *in-vitro* systems (waveguides at 900 and 1800 MHz; TEM cells, wire-patch cell and STUK resonator chamber), *in-vivo* (Mini Wheel, Mainly-Head and Circular Waveguide Setups). These systems were developed to provide technical quality control during the entire period of exposure. A thorough dosimetry was implemented including an analysis of possible artefacts, the continuous monitoring of exposure and environmental parameters and blinded exposure protocols. The three investigations have dealt with (i) genotoxicity, (ii) ODC activity in cells, and (iii) memory of rodents.

Using CW RF exposure at 935.2 MHz, Maes et al. had shown a weak increase in sister chromatid exchange in human lymphocytes consecutively treated with the genotoxic chemical mitomycin C (Maes et al., 1997). However, the Maes group was unable to replicate the data and thus a physical instead of chemical genotoxic agent was used in combination with RFR. Several standard *in vitro* tests for chromosomal and DNA damage in human lymphocytes were used. It was examined whether in these assays a 935 MHz GSM signal delivering SARs of 1 and 2 W/kg and a signal at 1800 MHz and 1 W/kg are genotoxic per se or can influence the genotoxicity of X-rays. Within the experiment parameters of the present study, in all instances no positive effect from the RFR signal was observed.

Ornithine decarboxylase, ODC, is a “marker enzyme” for cellular proliferation. The Litovitz research group has reported a temporary increase in ODC activity in L929 fibroblasts after exposure to RFR emitted from mobile phones (Penafiel et al., 1997). Only modulated RFR exhibited an effect on the ODC enzyme, compared to continuous wave signals (CW). All results of the Perform B programme did not support the previously reported effect of RFR on the activity of the ODC enzyme. The replication of the former work of the group of Litovitz, using 835 MHz/DAMPS and CW signals and murine L929 fibroblasts was extended to various RFR signals and to various cell types, including primary and live cells. Reminding that at least a ten-fold increase in ODC activity is necessary to induce deleterious effect on cellular functioning, the results give also no support for the possibility of neoplastic transformation via an effect on ODC activity.

There was limited animal evidence from the Lai group [Lai, et al. 1994] to suggest that pulse-modulated microwave radiation can transiently decrease radial arm maze performance in rodents (see also the previous chapter on memory). This is indicative of an effect on spatial working memory, a function associated in both rodents and humans with a part of the brain called the hippocampus. However, the results of the Perform B groups do not suggest that exposure to either pulsed 2.45 GHz fields or GSM-900 signals cause adverse effects on spatial

working memory in either rats or mice. Moreover, exposure to pulsed 2.45 GHz fields affects levels of behavioural anxiety in rats, or significantly affects the permeability of the BBB *in vivo*. In addition, similar negative results were obtained in mice whether they were exposed over the whole body or exposure was confined mainly to the head region. In particular, these experiments failed to confirm the results of studies performed by Lai and co-workers suggesting that exposure to RF fields induce large changes in maze behaviour in rodents. While these studies do not rule out possible more subtle effects on other aspects of cognition in animals, they strongly indicate that generalised deficits in behaviour are unlikely to occur.

## **REFLEX: Risk Evaluation of Potential Environmental Hazards from Low Energy Electromagnetic Field (EMF) Exposure Using Sensitive *in vitro* Methods**

The co-ordination of this European programme was done by the Verum Foundation in Munich. Twelve laboratories were involved in this highly structured multicentric study on ELF and RF field effects *in vitro*.

### *ELF magnetic fields*

Genotoxic effects were studied on primary cell cultures of human fibroblasts and on different cell lines. DNA strand breaks were observed in human fibroblasts at a significant level at 35  $\mu$ T [Ivancsits, et al. 2003a; Ivancsits, et al. 2003b; Ivancsits, et al. 2002]. The genotoxic effect was only observed when cells were exposed to intermittent fields and not to continuous exposure. The effect differed among the other types of cells examined. Confirmation studies in human fibroblasts in two other Reflex laboratories were carried out, with opposite results (one confirmed while the other did not). Chromosomal aberrations were also observed after exposure of human fibroblasts. Other observations were made in different laboratories: (i) ELF-EMF at around 2 mT upregulated the expression of early genes, such as p21, c-jun and egr-1, in p53-deficient mouse embryonic stem cells, but not in healthy wild-type cells [Czyz, et al. 2004b], (ii) ELF-EMF at 0.1 mT increased the proliferation rate of neuroblastoma cells, and (iii) ELF-EMF at 0.8 mT increased the differentiation of mouse stem cells into cardiomyocytes [Ventura, et al. 2004]. However, no effects were found on DNA synthesis, cell cycle, cell differentiation, cell proliferation and apoptosis in different cell types.

Thus, the few positive data on DNA damage originating from the Reflex programme appear to be in contradiction with results published in the literature. However, effects of intermittent ELF exposure have not yet been sufficiently investigated and therefore one must be careful in concluding about these data.

### *RF fields*

Some genotoxic effects were reported in fibroblasts, granulosa cells and HL60 cells. Cells responded to RF exposure between 0.3 and 2 W/kg with a significant increase in single and double strand DNA breaks and in micronuclei frequency (not published). Chromosomal aberrations in fibroblasts were also observed after exposure. Downregulation of the expression of neuronal genes in neuronal precursor cells and upregulation of the expression of early genes in p53-deficient embryonic stem cells were observed at 1.5 W/kg, but not in wild-type cells [Czyz, et al. 2004a]. Proteomic analyses on human endothelial cell lines showed that exposure altered the expression and phosphorylation of numerous, yet largely unidentified proteins [Leszczynski, et al. 2002; Leszczynski, et al. 2004; Nylund and Leszczynski 2004]. There was no evidence of effects on processes such as cell proliferation, apoptosis or immune cell functions [Capri, et al. 2004].

It is highly surprising that an effect of RF exposure could be present in a narrow range of SAR levels. One must thus await the publication in a scientific journal and replication of the positive data on DNA fragmentation which contradict all recent findings: a great majority of the published reports suggests that exposure of mammalian cells and animals does not result in increased DNA strand breaks, chromosomal aberrations, micronuclei and gene mutations (see Perform B above).

In conclusion, some of the positive data from the Reflex programme, when published, will warrant replication. However, research *in vivo* looking for effects possibly related to findings of the Reflex programme is premature, if not inappropriate.

# ***Sensitivity of children to EMF***

## **Introduction**

Children in both industrialized and developing countries are exposed to a large variety of chemical, physical and biological environmental exposures. Real or perceived increases in the incidence of certain childhood diseases, such as asthma, leukemia and brain cancer, and some behavioral and learning disabilities are being linked to environmental factors. Environmental exposures can be particularly harmful to children because of their special vulnerability during periods of development, and also because of the type and magnitude of exposure and the potential irreversibility of deleterious health effects. With rapid advances in technology resulting in larger numbers of sources exposing people around the world to higher levels of electromagnetic fields (EMF) over much of the electromagnetic spectrum, questions have been raised as to whether EMF might have particular effects on children.

With the rapid advances in EMF technologies and communications that are being used by children at earlier and earlier ages, thorough discussion is becoming urgent. To evaluate the available information relevant to children's sensitivity to both ELF- and RF-EMF and to identify research needs, the World Health Organization (WHO) held an expert workshop in Istanbul, Turkey in June 2004. This part of our report is based on discussions and recommendations from that workshop.

## **Development of children**

Embryos, infants, children and adolescents undergo different types of development. Development throughout the prenatal period is characterized by a highly ordered sequence of cell proliferation and differentiation, migration and programmed cell death (apoptosis). It is roughly divided into three periods: the preimplantation period, extending from fertilization to the settling of the embryo into the uterine wall; a period of organogenesis, characterized by the formation of the main body structures; and the fetal period, during which growth of the structures already formed takes place. The developing central nervous system seems to be particularly vulnerable because of the limited number and restricted physical location of the cohorts of proliferating cells from which it arises, and the precision required for building the complex neuronal architecture essential to proper brain function [Edwards, et al. 2003; Kimler 1998; Rodier 2004]. Major human tissue and organ development ends with the completion of sexual development at the end of the second or the beginning of the third decade of life. Development of the CNS continues through childhood and adolescence, largely through increased myelination and changes in synaptogenesis [Rodier 2004].

During the embryonic and fetal stages of development, exposure to environmental toxicants can result in death, congenital malformations, fetal or organ growth retardation, mental retardation, microcephaly, neurobehavioural effects, prematurity and stillbirth. During the postnatal stages of infant, child and adolescent development, exposure to toxic agents can result in total-organism growth retardation or selective tissue and organ retardation, cancer, interference with fertility and endocrine function, alterations in sexual maturation, interference with development of the immune system and alterations to neurological development. Of particular relevance to EMF exposure are childhood leukaemia and brain cancer and cognitive development.

Several aspects of exposure and susceptibility warrant a focus on children. In some exposure scenarios children may receive higher doses than adults owing to higher intake and accumulation or to differences in behaviour. Greater susceptibility to some toxicants and physical agents has been demonstrated in children, and although the mechanism for this increased susceptibility is not well understood, it is likely to involve greater cell division in growing and developing tissues. However, the actual increase in sensitivity is frequently contested, and there are even some examples of increased resistance. In addition, a longer expected lifetime, with a consequently increased chance of repeated exposure and accumulated damage, can also lead to higher risk, particularly for cancer. Some effects may not be clinically evident until long after exposure to a toxic agent

## **Dosimetry for children**

Numerous differences in the anatomy and physiology of children and adults are of particular consequence to dosimetry modelling for ELF and RF exposures. These parameters include differences in body shape and bone marrow distribution for ELF exposures; head size, skull thickness and ear elasticity for RF exposures; and dielectric properties for both.

In general, for ELF, adults exposed to low frequency electric or magnetic fields have higher internal electric field strengths and current densities than children because of size and shape differences. However, the distributions are different, and some tissues have higher current densities in children for the same external field. Furthermore, children have significantly higher internal field strengths and current densities from contact currents than do adults [Dawson, et al. 2001].

Modelling of SARs based on uniform or even nonuniform downscaling are inaccurate. To improve accuracy, MRI models, which take into account age-specific differences as well as variability between children, should be used. Despite large differences in the size, shape, and tissue distribution of heads, the SAR values and exposure variations for the child models are similar to those for adults, although somewhat higher. In addition, the relative depth of penetration is larger for children, a logical consequence of the smaller head diameter. Better modeling of the thickness of the compressed ears of children is needed. While some preliminary modeling indicated low exposure to fetuses, better models are being developed.

Dielectric studies encompassing several tissue types, including brain, obtained from newborn to fully grown rats, mice and rabbits exposed to RF-EMF in the frequency ranges of 130 MHz–10 GHz and 300 kHz to 300 MHz report large, age-related variations in the permittivity and conductivity of brain tissue, and even larger variations for skin and skull tissue [Peyman and Gabriel 2003; Peyman, et al. 2001; Thurai, et al. 1984; Thurai, et al. 1985]. Using age-related dielectric data to calculate whole-body and tissue-specific SARs in a scaled rat model led to significant differences in whole-body SARs. In addition, variations in tissue dielectric properties affect localized SAR values owing to corresponding variations of localized electric field and boundary conditions between neighboring tissues.

## **Exposure of children to RF and ELF Fields**

In evaluating the potential role of environmental exposures in the development of childhood diseases, it is important to consider not only the fact that childhood exposures can be different from exposures during adulthood, but also the fact that they can be highly age dependent. Exposures of interest during the preconception and gestation periods include residential and

parental exposures to ELF and RF fields as well as mothers' use of mobile phones, especially with a hands-free device. Infants and toddlers are exposed mostly at home or at day-care facilities. Among preteens, exposure sources expand to include mobile phone use and sources at school. Finally, adolescents are exposed to a wider variety of sources, including increasing use of mobile phones. Here we focus on two major exposure scenarios: residential ELF and RF exposures and exposure from mobile phones.

Everyone is exposed to ELF electric and magnetic fields at home. High-voltage power lines are a major source of exposure for children who live near them; however, only small proportion of children live in close proximity to high-voltage lines, in Sweden, for example, that number is less than one percent. There is evidence that younger children use appliances less (and spend less time outside the home), so that their personal exposure is closer to, and correlates better with, the field in the home.

RF fields are produced by radio and TV broadcasts, mobile phone base stations, and other communications infrastructure. Broadcast radio frequencies have been present for decades, but exposure to mobile phone signals has risen from zero in just a few years. While mobile phones are the dominant source of radiofrequency exposure for adolescents, environmental exposure to radio waves may be of concern for very young children. Note however, that exposure from base stations is orders of magnitude lower than that from mobile phones. The prevalence of mobile phone ownership among children is rising sharply and involves younger and younger children. In addition to the base stations, other sources such as TV and radio towers and deck phones at home contribute to the environmental RF exposure. Good information on population exposure and relative contributions of various sources is lacking. We consider it important for SSI to develop such information, which has become more feasible with the development of personal dosimeters.

When they reach adulthood, today's children will likely experience a longer period of exposure to RF fields from mobile phone and will have a much higher cumulative exposure than today's adults. However, rapidly changing technology might modify exposure. Radio-frequency exposure to children may be estimated more easily, because the variety of exposure sources is smaller for children than for adults.

## **Health effects**

The interaction of EMF with biological tissue may involve either electric or magnetic fields, both of which are known to produce effects at relatively high levels. At these levels, ELF fields act inducing currents in tissue, while RF fields produce heating. Mechanisms that could possibly lead to effects at low exposure levels remain speculative. The vast literature on in vivo and in vitro experiments of both ELF and RF fields using a variety of cell systems and species is not reviewed here. Rather, we focus on the mechanisms and experiments most relevant to the potential sensitivity of children.

## **Cancer in Children**

### **Extremely Low-Frequency Fields**

Most carcinogenesis bioassays begin when animals are sexually mature. A number of assays have been published showing the absence of initiating and promoting effects of ELF magnetic fields. Importantly, an animal model for B-cell acute lymphoblastic leukemia, the most common leukemia among children, was lacking until recently. The results of an ongoing

study of animals exposed to a 50-Hz magnetic field (100  $\mu\text{T}$ ) or a 50-Hz field plus harmonics using this newly developed model should be reported shortly [Bernard 2004]. These authors have found no evidence of an increase in incidence of leukemia in the exposed rats but have failed to provide positive controls showing that this animal model can be modulated. A number of *in vivo* studies have examined the effects of ELF fields on the responsiveness of the whole immune system. These studies have been carried out using differential white blood cell counts or standard *in vitro* tests of the immunological function of cells taken from the peripheral blood or spleen of exposed animals. The general consensus is that there is only weak evidence that power-frequency exposure can affect—and especially inhibit—the various aspects of immune system functions, including those relevant to cancer.

The potential risks to children of exposure to ELF-EMF, estimated from residential proximity to power sources and from measured fields, have been investigated in relation to *in utero* and postnatal time periods and to paternal exposure. Since an association between residential extremely low frequency electric and magnetic fields and childhood cancer, including leukemia, was first suggested in a report published in 1979, dozens of increasingly sophisticated studies have examined this association. Subsequently, a number of comprehensive reviews, several meta-analyses and two pooled analyses have summarized results from numerous sources. The two pooled analyses are closely consistent with each other and have reported the strongest associations between EMF and childhood leukemia in the literature: a risk of about two for children exposed to magnetic fields greater than 0.3–0.4  $\mu\text{T}$  compared with children exposed to fields less than 0.1  $\mu\text{T}$  [Ahlbom, et al. 2000; Greenland, et al. 2000]. Based on these results, IARC [IARC 2002] classified ELF magnetic fields as “possibly carcinogenic to humans.” While the results of the pooled analyses are not likely to be due to chance or to a known confounder, bias cannot be ruled out. The assessment of exposure to magnetic fields has improved over time, yet our ability to predict exposure remains severely limited—although it might be better for children than for adults [Forssén, et al. 2002]. An epidemiologically detectable risk of leukemia for children, but not for adults, might result from either better exposure assessment for children, or from greater susceptibility in children. Initially, it appeared that the EMF association was stronger with childhood brain tumors than with leukemia; however, several of the most recent studies have found no association [Kheifets 2001]. While no consistent associations have been observed for childhood CNS tumors, pooled analyses (proven so informative for childhood leukemia) of brain tumor studies have not been done.

### Radiofrequency Fields

Only a few studies have included prenatal or newborn rodent exposure. Adey et al. [Adey, et al. 1999] have exposed rats to American mobile telephone signals *in utero*. No effects were found on spontaneous tumorigenicity in the central nervous system in the offspring of exposed rats. Moreover, a trend for a decrease in the incidence of primary CNS tumors was shown for rats treated with a single dose of the neurocarcinogen ethylnitrosourea *in utero*. In other bioassays, animals were usually five or six weeks old and sexually mature (comparable to teenagers) at the start of a 2-year exposure period. In these bioassays, young animals were exposed for part of or the rest of their lives. Initiation as well as promotion was investigated. All of the studies have reported negative results in normal animals at SARs compatible with mobile telephony (e.g. [Anane, et al. 2003], while controversy still exists about the carcinogenic effects of RF radiation in a transgenic model [Repacholi, et al. 1997]. Major replications of this work should be completed soon. Gatta et al. [Gatta, et al. 2003] showed that immunological parameters were not altered in mice with whole-body exposure to RF fields at a SAR of 1 and 2 W/kg. Although cell differentiation plays a central role in the

development of organisms, only sparse information is available on the effects of RF radiation on differentiation. One study reported that exposure to RF fields at 1.5 W/kg up-regulated the expression of neuronal genes in neuronal precursor cells in p53-deficient embryonic stem cells, but not in wild-type cells [Czyz, et al. 2004a].

Several ecological studies [Cooper, et al. 2001; Dolk, et al. 1997a; Dolk, et al. 1997b; Hocking, et al. 1996; Maskarinec, et al. 1994; McKenzie, et al. 1998; Michelozzi, et al. 2002] have examined cancer risk, including risk of childhood leukemia, among populations living in proximity to radio and television broadcast towers. Driven by a previously identified cluster, these analyses are based simply on distance from the source and often include an extremely small number of cases. Such studies have been uninformative.

## **Developmental Effects and Pregnancy Outcomes**

### **Extremely Low Frequency**

Several studies, including a few with large group sizes and exposure over several generations, have evaluated exposure to electric fields of up to 150 kV/m in several mammalian species. The results of these studies are rather consistent, and do not suggest adverse developmental effects [Juutilainen 2003]. Magnetic field exposure has been evaluated both in non-mammalian and mammalian models. Results from several independent research groups suggest that exposure to ELF magnetic fields at microtesla levels may disturb early development of bird embryos. However, replication attempts have been unsuccessful in some laboratories. Results from experiments with other non-mammalian experimental models (fish, sea urchins and insects) have also suggested subtle effects on developmental stability. In mammals, prenatal exposure to ELF magnetic fields does not result in strong adverse effects on development. Gross external, visceral or skeletal malformations are not increased by fields of up to 20 mT (at 50 Hz). The only finding that shows some consistency is an increase in minor skeleton alterations in several experiments, in both rats and mice. Skeletal variations are relatively common findings in teratological studies and are often considered biologically insignificant. Nevertheless, subtle effects of magnetic field exposure on skeletal development cannot be ruled out. Some effects of magnetic (or combined electric and magnetic) fields on postnatal development have been reported, but evaluation of the consistency of the findings is difficult owing to the varying methods and approaches used in different studies.

Numerous epidemiological studies of various pregnancy outcomes in relation to EMF are available in the scientific literature. They include studies investigating the use of video display terminals (VDTs), electric blankets or heated waterbeds, as well as studies of parental occupational exposure. Most studies have found no effects, but these studies are limited in exposure assessment and lack the power to examine high exposure levels. Two studies that have included personal measurements of ELF-EMF exposure have reported effects on spontaneous abortion in relation to maximum measured magnetic fields [Lee, et al. 2002; Li, et al. 2002]. The possibility of exposure assessment bias in these studies has been discussed, and results need to be confirmed in additional studies before firm conclusions can be drawn.

### **Radiofrequency Fields**

Numerous studies have evaluated developmental effects of RF fields on mammals, birds and other non-mammalian species [Heynick and Merritt 2003]. These studies have shown clearly that RF fields are teratogenic at exposure levels that are high enough to cause significant increases in temperature. There is no consistent evidence of effects at nonthermal exposure levels.

Several studies of occupational RF exposure, primarily to physiotherapists, have reported an increased risk of congenital malformations. However, no specific type of malformation has been consistently reported, and there is a potential for recall bias in these studies. Exposure to the fetus from a mobile phone kept in a pocket, handbag, or belt by the hip when a pregnant woman is using hands-free equipment has been mentioned. So far, no studies are available on pregnancy outcomes related to mobile telephony.

## **Behavioural and Cognitive Effects in Children**

Effects on brain function that lead to detrimental effects on cognition and behavior can be due to the high sensitivity of the developing and maturing nervous system to particular physical and chemical agents. Very few laboratory studies of the effects of RF fields and no studies of the effects of ELF fields on children appear to have been performed. Extrapolation of acute effects is necessary, using results from studies of adult volunteers and studies of both immature and adult animals. Although such extrapolation may provide some qualitative information regarding potential outcomes in children, it may not provide reliable estimates of risk.

### **Extremely Low-Frequency Fields**

The results of acute exposure to magnetic fields on simple and choice reaction time in adult volunteers do not suggest any field-dependent effects, although modest changes in speed and accuracy during task performance have been reported. These data also suggest that effects may depend on the difficulty of the task performed. A few studies have reported subtle field-dependent changes in other cognitive functions, including the performance of specific memory and attention tasks. Studies measuring the electrical activity of the brain have reported scattered field-dependent effects and hints of subtle trends, but no well-defined, field-dependent results.

Various animal models have been used to investigate possible field-induced effects on brain function, electrical activity and behavior. These effects include changes in neurotransmitter levels, electrical activity, and the performance of learned tasks. Overall, while a few field-dependent responses, such as changes in spatial memory, have been tentatively identified, the fragmented nature of the observed responses does not suggest an obvious deleterious effect. The available evidence is thus suggestive of subtle effects, but is not conclusive.

### **Radiofrequency Fields**

Several recent studies have examined the effects of RF fields associated with mobile phones on attention, memory and other cognitive functions. No consistent responses have been observed, and these studies provide only weak evidence that changes in cognitive performance occur following low-level exposure. A few recent experimental studies have investigated the effects of exposure from Global System for Mobile Communications (GSM) signals at 902 MHz on child cognition [Preece unpublished]. Finnish researchers have recently investigated the effects of mobile phones on brain activity, cognitive functions and subjective symptoms in 10-14 year old children. Peer-reviewed publications are not yet available but according to a preliminary report [Haarala and Krause 2003], some effects were seen on EEG of children performing an auditory memory task, while no effects were observed on cognitive functions or subjective symptoms). In the other study, mobile phone exposure did not affect cognitive performance as measured by response speed and accuracy. One observational study found a mild facilitating effect on attention, as measured in paper and

pencil tests, in adolescent users of mobile phones, but this may reflect changes in motor skills rather than attention per se [Lee, et al. 2001]. Several studies have reported that RF fields may affect the electrical activity of the brain. However, no specific response has been identified, and other studies have not reported any changes. Some, but not all, studies suggest that RF fields may affect certain components of sleep.

The possible effects of RF fields in animals have been investigated using a variety of approaches, from measurement of specific gene expression and neurotransmitter activity to investigation of changes in learned behaviors and in the electrical activity of the brain. The majority of reported RF effects appear to be consistent with hyperthermia or thermoregulation, or with stresses associated with exposure. Some data suggest that spatial maze behavior of rats can be impaired by exposure to pulsed RF fields in the absence of whole-body heating, although more recent studies found no significant RF effects. In addition, two studies indicate that the learned behavioral responses of adult rats were not affected by prenatal or early postnatal exposure to low-level RF fields.

### **Other Outcomes**

Nothing is known about potential adverse health effects later in life from EMF (ELF and RF) exposures during childhood. Associations between occupational ELF magnetic field exposure and amyotrophic lateral sclerosis (ALS) and Alzheimer's disease have been reported, although alternative explanations to the findings are being discussed. There are also controversial reports that exposure to mobile-phone frequency fields affects the permeability of the blood-brain barrier. Perhaps we need to ask whether EMF exposure in childhood can increase the risk of developing Alzheimer's disease, brain tumors, or other outcomes later in life, especially in light of extensive mobile phone use among teenagers.

### **Research gaps**

Physical, chemical and therapeutic agents have the potential for affecting development, depending on the nature of the agent and the timing and magnitude of the exposure. For ELF magnetic fields there are indications that children might be more sensitive; however, we lack the understanding of how, or even if, these fields might be involved in leukemogenesis. Several analyses can help advance the current state of knowledge, particularly on the potential effect of ELF on children. These include update of the pooled analysis of childhood leukemia, pooled analysis of brain cancer studies, studies of large highly exposed and/or susceptible populations, evaluation of selection bias and other hypothesis such as contact current and melatonin. Meanwhile, given the current state of science and despite the lack of a known mechanism precautionary approach is warranted.

Even less is known about the potential sensitivity of children to RF fields. Widespread exposure to these fields is recent, and although much of the laboratory research has been reassuring, its focus has not been on children. Moreover, few epidemiologic studies of adequate quality have been completed and even fewer are of relevance to children. Given the paucity of data indicating a particular vulnerability of children to EMF, it may be tempting to conclude that children are not more susceptible than adults to RF exposure. However, the absence of an observed effect does not necessarily mean that exposure is harmless, especially if crucial studies focusing on children are yet to be done. In addition, many of the methods used have not been sensitive enough to detect subtle effects. In the absence of direct evidence for or against the potential susceptibility of children to EMF, one can extrapolate either from

data on adults or from data on other chemical and physical agents. Extrapolation from data on adults is problematic. Because the period from embryonic life to adolescence is characterized by growth and development, deleterious effects can occur at lower levels be more severe, or lead to effects that do not occur in adults; on the other hand, children can be more resilient, owing to better recuperative capacities. Extrapolating from other exposures is also problematic: while it is clear that children are more vulnerable to many physical and chemical agents, many of their effects are very specific in terms of both the level and timing of exposure. Thus, there is an urgent need to design and implement studies that can provide direct and relevant information on the potential susceptibility of children to RF exposure. Owing to widespread use of mobile phones and relatively high exposures to the brain among children and adolescents, investigation of the potential effects of RF fields on cognition See [www.who.int/peh-emf/research/rf03/en](http://www.who.int/peh-emf/research/rf03/en) for more details on a research agenda that identifies high-priority studies needed to fully assess the potential vulnerability of children to ELF and RF fields and outlines the rationale for these studies.

### **Limited scientific evidence on the carcinogenicity of electric and magnetic fields - is it enough for precautionary preventive action?**

Evaluating the risk related to various agents with potential to increase cancer risk in humans is often controversial. Several national and international health agencies have established programs with the aim of identifying carcinogens.

In principle, it is an activity grounded in the scientific evaluation of the results of human epidemiological studies, long-term bioassays in experimental animals and other data relevant to an evaluation of carcinogenicity and its mechanisms. In this brief commentary, we discuss the interpretation of the categories adopted in the IARC Monograph program to identify carcinogens.

The IARC Monographs are using the following descriptors:

**Table 1: IARC Categories of evidence**

<b>Categories</b>	<b>Cancer hazard to human</b>
<b>Group 1</b>	the agent or mixed exposure is carcinogenic to humans
<b>Group 2 A</b>	the a the agent or mixed exposure is probably carcinogenic to humans
<b>Group 2 B</b>	the agent or mixed exposure is possibly carcinogenic to humans
<b>Group 3</b>	the agent or mixed exposure is not classifiable as to its carcinogenicity
<b>Group 4</b>	the agent or mixed exposure is probably not carcinogenic to humans

#### **Epidemiological data in humans**

Assignments to Group 1, human carcinogens, are made exclusively on the basis of human data, the experimental data playing no role (see Table 2). However, there are a few notorious cases, where an agent has been classified into the group in the absence of sufficient evidence of carcinogenicity in humans. An example is TCDD. This was done based on convincing animal data and good understanding of the mechanistic principles of cancer causation in

rodents. There was enough evidence in humans which indicated similar mechanisms and also operative in humans, making the probability that TCDD would also be carcinogenic to humans high.

**Table 2: Overall default evaluation of carcinogenicity in the IARC Monographs**

		Cancer in experimental animals		
		Sufficient	Limited	Inadequate
Cancer in humans	Sufficient	1 Carcinogenic	1 Carcinogenic	1 Carcinogenic
	Limited	2A Probably carcinogenic	2B Possibly carcinogenic	2B Possibly carcinogenic
	Inadequate	2B Possibly carcinogenic	3 Not known	3 Not known

In making the assessment of causation in humans, epidemiologists ask whether a causal association is credible and whether chance, bias, or confounding factors can be excluded. The guidance from the set of "Hill criteria" is searched, including

- Consistency of the observed association
- Strength of the observed association
- Specificity of the observed association
- Temporal relationship of the observed association
- Biological gradient (exposure-response relationship)
- Biological plausibility
- Coherence
- Experimental evidence (from human populations)
- Analogy

### Experimental Animal Bioassay Data

The most common way of identifying potentially carcinogenic agents is long-term bioassays in experimental animals. Experimental carcinogenesis research is based on the assumption that agents, which can cause cancer in animals, will have similar effects in humans. Within the IARC Monographs program, the respective roles of human and experimental data in defining and/or predicting human risk have been amply and repeatedly debated. It has been agreed that carcinogenic risk to humans could be quantified on the basis of experimental animal data only with large uncertainties. However, experimental evidence has maintained its validity in demonstrating carcinogenicity of an agent and in serving as an alert that similar effects, might occur in humans. Accordingly, the IARC monographs have a policy statement in the preamble, which states that in the absence of adequate human data, it is biologically plausible and prudent to regard agents and mixtures for which there is sufficient evidence of carcinogenicity in experimental animals as if they presented a carcinogenic risk to humans [IARC 2002]. In doing so, IARC, anticipating what later became known as the 'precautionary principle', was trying to reconcile a scientifically accurate analysis of the data with an interpretation of the evidence of carcinogenicity provided by experimental data that is not

only biologically plausible but is public health orientated and gives priority to primary prevention.

### Evaluation of the data on mechanisms

Consideration of the mechanistic data has the potential of improving the analysis of causation between the observed effect and the given exposure. Elucidation of the mechanisms gives insight into the potential steps important in the process and prevention of carcinogenesis. There may be strong associations between exposure and disease, but if confounding factors are not examined thoroughly, the association may be spurious.

### The IARC categories and need for cancer prevention

The IARC categories are considered to be "hazard" assessments, not risk assessments, which would include an attempt for quantitation of the potential risk. This risk depends on both the existence of hazard and exposure to that hazard. A cancer hazard exists when an agent has been categorized to Groups 1, 2A and 2B (see Table 1). It should be emphasized that the categorization of an agent with regard to its carcinogenicity remains a matter of scientific judgment, and agents or exposures can be classified to different categories in a way different from the strict interpretation of the default criteria. In Group 1 the hazard has been shown to produce a real risk in humans (as observed in epidemiological studies of exposed humans). In Group 2A there are clear indications of the cancer hazard, and in Group 2B the hazard is considered to be high enough to consider, for practical purposes, preventive actions. The agents which cause cancer in animal models, but do not have any relevant human data, are categorized in Group 2B as 'possible human carcinogens'.

The category 2B is generally used for agents for which there is limited evidence of carcinogenicity in humans in the absence of sufficient evidence of carcinogenicity in experimental animals, or when there is inadequate evidence of carcinogenicity in humans but there is sufficient evidence of carcinogenicity in experimental animals. The above mentioned public health statement in the IARC Monographs preamble that agents with 'sufficient evidence' in experimental animal models should be considered, for practical purposes and for the purpose of primary prevention, 'as if they presented a carcinogenic risk to humans' should logically also apply to all agents in this category 2B of 'possible human carcinogens'. The preventive approaches to be considered could include strategies to reduce exposures as much as technologically possible, strategies to search for alternative chemicals/agents etc.

### Evaluation of carcinogenicity of electric and magnetic fields in the IARC Monographs

Exposure to electric and magnetic fields (EMF) emanating from utilization of electricity is widespread. The major concern has focused on the possibility that exposure to extremely low-frequency (ELF) electric and magnetic fields may result in increased risk of cancer. It is a widely held view that the current scientific evidence in support of an association between ELF magnetic fields and childhood leukemia is limited [IARC 2002]. The evaluation in the IARC monographs lead to overall evaluation of Group 2B, ELF magnetic fields being possibly carcinogenic to humans. Evidence for static electric and magnetic fields and ELF electric fields is not classifiable as to their carcinogenicity to humans (Group 3). As far as radiofrequency fields (RF) are concerned, the evaluation in the IARC Monographs has not yet been done, and it is planned to take place in 2006.

## Developing Policy for Children

Given the paucity of data indicating a particular vulnerability of children to EMF, there is an urgent need to develop policies in the face of uncertainty and to design and implement studies that can provide direct and relevant information on the potential susceptibility of children.

### Setting Guidelines

Guideline-setting bodies often recognize that children might be more susceptible to EMF than adults, that EMF dosimetry is different in children and that behavior, and thus exposure, might be different in children. The potential for these differences is incorporated in an additional reduction factor that is applied to exposure limits for the general population (as compared to limits for occupational exposure), rather than the development of specific limits for children.

The exposure guidelines developed by the International Commission on Non-ionizing Radiation Protection (ICNIRP) are intended to protect against adverse health effects that are scientifically established. In the case of electromagnetic fields, all established effects are acute ones [ICNIRP 2002] with specific thresholds, in the sense that a minimum biologically effective quantity must be applied for an effect to occur. In this approach, the concept of *critical effect* is fundamental. The critical effect is the established adverse health effect that occurs at the lowest level of exposure. If a given effect were found to occur in children—or in any other particular group—at an exposure level lower than that required for an effect in the rest of the population, it would be assumed to be critical. Sources of uncertainty include possible differences between children and adults in the distribution of induced currents and in SARs (for low- and high-frequency fields, respectively). The reduction factor of 5 used for both ELF and RF exposures for the general population is likely to be adequate for the protection of children against established health effects.

### Precautionary Approaches

The precautionary principle is an attempt to link the scientific community and the community of 'common' people in joint responsibility for decisions, a state where primary importance attaches to our right to be informed and our freedom of choice. Risk involved in electromagnetic fields may well be lower than in many other areas currently well accepted (such as risks of motor traffic), but public opinion bluntens the perception of certain risks, as in case of ultrafine particles from motor exhausts. Scientific evidence can make its contribution to societal decision making openly, to clarify the contours of uncertainty with the reservations typical of scientific rationality. For its part, the political decision making procedure must supply the tools and channels for democracy to find its voice.

While development of science-based guidelines for established effects is challenging, development of science-based precautionary approaches for uncertain effects is even more difficult, particularly when children may be affected. Many societies have a heightened level of concern for vulnerable populations, particularly children, because they may be unable to take actions to effectively manage their own risk. Furthermore, many societies believe that children and fetuses should be afforded an even higher level of protection because they represent the future of the society (WHO Precautionary Framework, 2004). Other factors, such as the potential number of productive years lost and the potential for higher and longer

exposure, also come into consideration. Finally, widespread exposure can have large public health consequences even if its effects are small or subtle.

The draft WHO Precautionary Framework discusses several factors that argue for the adoption of greater, rather than lesser, protection for EMF exposure to children. Among these factors are the possibility that EMF might affect children; the dread with which some of the diseases in question, such as leukemia and brain cancer, are perceived; the involuntary nature of some of the exposure; the extensiveness of exposure; the rapid growth of some exposures; and the occurrence of some of the exposures to children close to guideline limits. Possible precautionary measures for EMF will vary from country to country, but should follow WHO Precautionary Framework recommendations as outlined in the case studies. In Sweden, such approaches have already been introduced for both ELF and RF fields.

The precautionary principle emerged as a decision rule for regulating environmentally hazardous activities in the Swedish Environmental Protection Act of 1969. This act which remains in effect today incorporates the statement that the mere risk of harm, if not remote, warrants protective measures or a ban on the activity that is possibly causing harm. For low frequency fields, the Swedish Radiation Protection Authority recommended reducing exposures specifically without providing any recommendation regarding levels. In 1993 the Swedish government, acting upon the recommendation of the Swedish Electric Board [Villa and Ljung 1993], advocated prudent avoidance. This cautionary policy, defined as taking measures to reduce magnetic fields in newly built housing and electrical facilities without great inconvenience or cost, was formalized in 1996 in a guide for Swedish local officials [National Board of Occupational Safety and Health, et al. 1996]: "If measures generally reducing exposure can be taken at reasonable expense and with reasonable consequences in all other respects, an effort should be made to reduce fields radically deviating from what could be deemed normal in the environment concerned. Where new electrical installations and buildings are concerned, efforts should be made to design and position them in such a way that exposure is limited."

For RF, the joint statement of Scandinavian Radiation Protection last year adopted precaution (2004)

"The existing knowledge gaps and the prevailing scientific uncertainty justify a certain precautionary attitude regarding the use of handsets for mobile telephony. Due to the widespread use of mobile phones even a very small risk could have consequences for public health. Because of the lack of knowledge in certain fields of research the Nordic authorities find it is wise to use, for instance, a hands-free kit that reduces the exposure to the head significantly. This information should be addressed both to adults, young people and children. It is important that parents inform young people and children about how to reduce the exposure from mobile phones."

## ***Discussion***

It is clear from the presentations and discussions in this report that there are several important ongoing studies and that there are several unresolved issues. Most of the current focus is on RF rather than on ELF or other lower frequencies. The most likely explanation to this is that ELF research began earlier and has come further. At least for the epidemiologic studies it may be reasonable to compare the current RF epidemiology to ELF epidemiology about a decade ago. A major explanation to this is that knowledge about exposure distribution in the population and measurement technology has developed further and is now much better for ELF fields. However, exposure assessment was a major obstacle in earlier ELF epidemiology and it remains so in current RF epidemiology. A number of second generation RF studies are in the process of being planned or piloted. A major issue in this process is the exposure assessment.

To date, little is known about the levels of radiofrequency exposure in the general population from sources such as mobile phones being used by oneself or other people, mobile phone base stations, and radio and television transmitters. Measurements that have been performed have usually been made as a result of public concern about base station exposures or other specific sources, and have therefore been made at locations that could be assumed to have higher fields than would be the case if measurement locations were selected randomly. Furthermore, all measurements have been stationary, and there is today no knowledge about the level of exposure that an individual will have throughout the day. Some countries have set up networks monitoring the radiofrequency exposure at certain locations, but again, the locations chosen have been driven by the existence of base stations or other RF exposure sources, and do not reflect the RF exposure in the general population.

There is a need for information about the personal exposure to RF fields in the general population, to enhance the understanding of the relative importance of exposure from base stations close to the home, from radio and television transmitters, and from the use of mobile phones. There is now a personal meter available that can measure the RF fields over a longer time period, e.g. 24 hours, and that has the possibility to distinguish between different sources of RF exposure. Studies with personal RF exposure measurements of randomly selected samples of the general population are strongly encouraged.

## References

- Adey WR, Byus CV, Cain CD, Higgins RJ, Jones RA, Kean CJ, Kuster N, MacMurray A, Stagg RB, Zimmerman G and others. 1999. Spontaneous and nitrosourea-induced primary tumors of the central nervous system in Fischer 344 rats chronically exposed to 836 MHz modulated microwaves. *Radiat Res* 152(3):293-302.
- Ahlbom A, Day N, Feychting M, Roman E, Skinner J, Dockerty J, Linet M, McBride M, Michaelis J, Olsen JH and others. 2000. A pooled analysis of magnetic fields and childhood leukaemia. *Br J Cancer* 83(5):692-8.
- Anane R, Dulou PE, Taxile M, Geffard M, Crespeau FL, Veyret B. 2003. Effects of GSM-900 microwaves on DMBA-induced mammary gland tumors in female Sprague-Dawley rats. *Radiat Res* 160(4):492-7.
- Andersson B, Berg M, Arnetz BB, Melin L, Langlet I, Liden S. 1996. A cognitive-behavioral treatment of patients suffering from "electric hypersensitivity". Subjective effects and reactions in a double-blind provocation study. *J Occup Environ Med* 38(8):752-8.
- Auvinen A, Hietanen M, Luukkonen R, Koskela RS. 2002. Brain tumors and salivary gland cancers among cellular telephone users. *Epidemiology* 13(3):356-9.
- Bergqvist U, Hillert L, Birke E. 2000. Elöverkänslighet och hälsorisker av elektriska och magnetiska fält. *Forskningsöversikt och utvärdering (In Swedish)*. arbetslivsforskning Rf, editor. Katrineholm: Hellmans Förlag.
- Bergqvist U, Vogel E, Aringer L, Cunningham J, Fabriziomaria G, Leitgeb N, Miro L, Neubauer G, Ruppe I, Veccia P and others. 1997. Possible health implications of subjective symptoms and electromagnetic fields. A report prepared by a European group of experts for the European Commission, DG V. Arbetslivsinstitutet, editor. Solna: Arbetslivsinstitutet.
- Bernard N. 2004. Mise au point d'un modèle de leucémie aiguë lymphoblastique de type B, chimiquement induite chez le rat pour l'étude d'agents leucémogènes : application aux champs magnétiques de 50 Hz. [Design of an ALL model of the B type, chemically-induced in the rat for the study of leukemic agents: application to 50-Hz magnetic fields]: University of Paris.
- Capri M, Scarcella E, Bianchi E, Fumelli C, Mesirca P, Agostini C, Remondini D, Schuderer J, Kuster N, Franceschi C and others. 2004. 1800 MHz radiofrequency (mobile phones, different Global System for Mobile communication modulations) does not affect apoptosis and heat shock protein 70 level in peripheral blood mononuclear cells from young and old donors. *Int J Radiat Biol* 80(6):389-97.
- Cassel J-C, Cosquer B, Galani R, Kuster N. 2004. Whole-body exposure to 2.45 GHz electromagnetic fields does not alter radial-maze performance in rats. *Behav Brain Res* 155:37-43.
- Chia SE, Chia HP, Tan JS. 2000. Prevalence of headache among handheld cellular telephone users in Singapore: a community study. *Environ Health Perspect* 108(11):1059-62.
- Christensen HC, Schuz J, Kosteljanetz M, Poulsen HS, Thomsen J, Johansen C. 2004. Cellular telephone use and risk of acoustic neuroma. *Am J Epidemiol* 159(3):277-83.
- Cobb B, Jauchem J, Adair E. 2004. Radial arm maze performance of rats following repeated low level microwave radiation exposure. *Bioelectromagnetics* 25:49-57.
- Cook C, Thomas A, Prato F. 2002. Human electrophysiological and cognitive effects of exposure to ELF magnetic and ELF modulated RF and microwave fields: a review of recent studies. *Bioelectromagnetics* 23:144-157.

- Cooper D, Hemmings K, Saunders P. 2001. Re: "Cancer incidence near radio and television transmitters in Great Britain. I. Sutton Coldfield transmitter; II. All high power transmitters". *Am J Epidemiol* 153(2):202-4.
- Cosquer B, Galani R, Kuster N, Cassel J-C. 2004. Whole-body exposure to 2.45 GHz electromagnetic fields does not alter anxiety responses in rats: A plus-maze study including test validation. *Behav Brain Res*(in press).
- Crasson M. 2003. 50-60 Hz electric and magnetic field effects on cognitive function in humans: a review. *Radiat Prot Dosimetry* 106(4):333-40.
- Curcio G, Ferrara M, De Gennaro L, Cristiani R, D'Inzeo G, Bertini M. 2004. Time-course of electromagnetic field effects on human performance and tympanic temperature. *Neuroreport* 15(1):161-4.
- Czyz J, Guan K, Zeng Q, Nikolova T, Meister A, Schönborn F, Schuderer J, Kuster N, AM W. 2004a. High frequency electromagnetic fields affect gene expression levels in tumor suppressor p53-deficient embryonic stem cells. *Bioelectromagnetics* 25:296-307.
- Czyz J, Nikolova T, Schuderer J, Kuster N, Wobus AM. 2004b. Non-thermal effects of power-line magnetic fields (50 Hz) on gene expression levels of pluripotent embryonic stem cells-the role of tumour suppressor p53. *Mutat Res* 557(1):63-74.
- D'Andrea J, Chou C, Johnston S, Adair E. 2003. Microwave effects on the nervous system. *Bioelectromagnetics Suppl* 6:S:107-147.
- Dawson TW, Caputa K, Stuchly MA, Kavet R. 2001. Electric fields in the human body resulting from 60-Hz contact currents. *IEEE Trans Biomed Eng* 48(9):1020-6.
- D'Costa H, Trueman G, Tang L, Abdel-rahman U, Abdel-rahman W, Ong K, Cosic I. 2003. Human brain wave activity during exposure to radiofrequency field emissions from mobile phones. *Australas Phys Eng Sci Med* 26:162-167.
- Dolk H, Elliott P, Shaddick G, Walls P, Thakrar B. 1997a. Cancer incidence near radio and television transmitters in Great Britain. II. All high power transmitters. *Am J Epidemiol* 145(1):10-7.
- Dolk H, Shaddick G, Walls P, Grundy C, Thakrar B, Kleinschmidt I, Elliott P. 1997b. Cancer incidence near radio and television transmitters in Great Britain. I. Sutton Coldfield transmitter. *Am J Epidemiol* 145(1):1-9.
- Dreyer NA, Loughlin JE, Rothman KJ. 1999. Cause-specific mortality in cellular telephone users. *Jama* 282(19):1814-6.
- Dubreuil D, Jay T, Edeline J-M. 2002. Does head-only exposure to GSM-900 MHz electromagnetic fields affect the performance of rats in spatial learning task? *Behav Brain Res* 129:203-210.
- Dubreuil D, Jay TM, Edeline J-M. 2003. Head-only exposure to GSM-900 MHz electromagnetic fields does not alter rat's memory in spatial and non-spatial tasks. *Behav Brain Res* 145:51-61.
- Edelstyn N, Oldershaw A. 2002. The acute effects of exposure to the electromagnetic field emitted by mobile phones on human attention. *Neuroreport* 13:119-121.
- Edwards MJ, Saunders RD, Shiota K. 2003. Effects of heat on embryos and foetuses. *Int J Hyperthermia* 19(3):295-324.
- Flodin U, Seneby A, Tegenfeldt C. 2000. Provocation of electric hypersensitivity under everyday conditions. *Scand J Work Environ Health* 26(2):93-8.
- Forssén UM, Ahlbom A, Feychting M. 2002. Relative contribution of residential and occupational magnetic field exposure over twenty-four hours among people living close to and far from a power line. *Bioelectromagnetics* 23(3):239-44.

- Gatta L, Pinto R, Ubaldi V, Pace L, Galloni P, Lovisolo GA, Marino C, Pioli C. 2003. Effects of in vivo exposure to GSM-modulated 900 MHz radiation on mouse peripheral lymphocytes. *Radiat Res* 160(5):600-5.
- Greenland S, Sheppard AR, Kaune WT, Poole C, Kelsh MA. 2000. A pooled analysis of magnetic fields, wire codes, and childhood leukemia. Childhood Leukemia-EMF Study Group. *Epidemiology* 11(6):624-34.
- Haarala C, Bjornberg L, Ek M, Laine M, Revonsuo A, Koivisto M, Hämäläinen H. 2003. Effect of a 902 MHz electromagnetic field emitted by mobile phones on human cognitive function: A replication study. *Bioelectromagnetics* 24:283-288.
- Haarala C, Krause C. 2003. Effects of mobile phones on human brain- and cognitive functions, and subjective symptoms. In: Juutilainen J, Kumlin T, editors. *Mobile telephony and health. Proceedings, Final seminar of the Finnish national research programme 1998-2003*. Helsinki, 17 October 2003.
- Hamblin D, Wood A. 2002. Effects of mobile phone emissions on human brain activity and sleep variables. *Int J Radiat Biol* 78:659-669.
- Hardell L, Hallquist A, Mild KH, Carlberg M, Pahlson A, Lilja A. 2002. Cellular and cordless telephones and the risk for brain tumours. *Eur J Cancer Prev* 11(4):377-86.
- Hardell L, Mild KH, Pahlson A, Hallquist A. 2001. Ionizing radiation, cellular telephones and the risk for brain tumours. *Eur J Cancer Prev* 10(6):523-9.
- Heynick LN, Merritt JH. 2003. Radiofrequency fields and teratogenesis. *Bioelectromagnetics Suppl* 6:S174-86.
- Hietanen M, Hamalainen AM, Husman T. 2002. Hypersensitivity symptoms associated with exposure to cellular telephones: no causal link. *Bioelectromagnetics* 23(4):264-70.
- Hillert L, Berglind N, Arnetz BB, Bellander T. 2002. Prevalence of self-reported hypersensitivity to electric or magnetic fields in a population-based questionnaire survey. *Scand J Work Environ Health* 28(1):33-41.
- Hinrikus H, Parts M, Lass J, Tuulik V. 2004. Changes in human EEG caused by low level modulated microwave stimulation. *Bioelectromagnetics* 25:431-440.
- Hocking B, Gordon IR, Grain HL, Hatfield GE. 1996. Cancer incidence and mortality and proximity to TV towers. *Med J Aust* 165(11-12):601-5.
- Huber R, Schuderer J, Graf T, Jutz K, Borbely A, Kuster N, Achermann P. 2003. Radio frequency electromagnetic field exposure in humans: Estimation of SAR distribution in the brain, effects on sleep and heart rate. *Bioelectromagnetics*. 24:262-276.
- Huber R, Treyer V, Borbely A, Schuderer J, Gottselig J, Landolt H, Werth E, Berthold T, Kuster N, Buck A and others. 2002. Electromagnetic fields, such as those from mobile phones, alter regional cerebral blood flow and sleep and waking EEG. *J Sleep Res*. 11:289-995.
- IARC. 2002. Non-ionizing radiation. Part 1, static and extremely low-frequency (ELF) electric and magnetic fields. Vol 80, Lyon.
- ICNIRP. 2002. General approach to protection against non-ionizing radiation. *Health Phys* 82(4):540-8.
- IEGMP. 2000. Independent Expert Group on Mobile Phones (Chairman: Sir William Stewart). *Mobile phones and health*. Chilton, Didcot, available at: <http://www.iegmp.org.uk/>.
- Inskip PD, Tarone RE, Hatch EE, Wilcosky TC, Shapiro WR, Selker RG, Fine HA, Black PM, Loeffler JS, Linet MS. 2001. Cellular-telephone use and brain tumors. *N Engl J Med* 344(2):79-86.
- Ivancsits S, Diem E, Jahn O, Rüdiger H. 2003a. Age-related effects on induction of DNA strand breaks by intermittent exposure to electromagnetic fields. *Mech Age Dev*.

- Ivancsits S, Diem E, Jahn O, Rüdiger H. 2003b. Intermittent extremely low frequency electromagnetic fields cause DNA damage in a dose dependent way. *Int Arch Occup Environ Health* 76:431-436.
- Ivancsits S, Diem E, Rüdiger H, Jahn O. 2002. Induction of DNA strand breaks by exposure to extremely-low-frequency electromagnetic fields in human diploid fibroblasts. *Mutation Res* 519:1-13.
- Johansen C, Boice J, Jr., McLaughlin J, Olsen J. 2001. Cellular telephones and cancer--a nationwide cohort study in Denmark. *J Natl Cancer Inst* 93(3):203-7.
- Juutilainen J. 2003. Developmental effects of extremely low frequency electric and magnetic fields. *Radiat Prot Dosimetry* 106(4):385-90.
- Kheifets LI. 2001. Electric and magnetic field exposure and brain cancer: a review. *Bioelectromagnetics Suppl* 5:S120-31.
- Kimler BF. 1998. Prenatal irradiation: a major concern for the developing brain. *Int J Radiat Biol* 73(4):423-34.
- Koivisto M, Haarala C, Krause CM, Revonsuo A, Laine M, Hamalainen H. 2001. GSM phone signal does not produce subjective symptoms. *Bioelectromagnetics* 22(3):212-5.
- Koivisto M, Krause C, Revonsuo A, Laine M, Hamalainen H. 2000. The effects of electromagnetic field emitted by GSM phones on working memory. *Neuroreport* 11:1641-1643.
- Kramarenko A, Tan U. 2003. Effects of high-frequency electromagnetic fields on human EEG: a brain mapping study. *Int J Neurosci* 113:1007-1019.
- Krause C, Haarala C, Sillanmaki L, Koivisto M, Alanko K, Revonsuo A, Laine M, Hamalainen H. 2004. Effects of electromagnetic field emitted by cellular phones on the EEG during an auditory memory task: a double blind replication study. *Bioelectromagnetics* 25:33-40.
- Kurokawa Y, Nitta H, Imai H, Kabuto M. 2003. No influence of short-term exposure to 50-Hz magnetic fields on cognitive performance function in human. *Int Arch Occup Environ Health* 76(6):437-42.
- Lai H. 2004. Interaction of microwaves and a temporally incoherent magnetic field on spatial learning in the rat. *Physiol Behav* 82:785-789.
- Lai H, Horita A, Guy A. 1994. Microwave irradiation affects radial-arm maze performance in the rat. *Bioelectromagnetics* 15:95-104.
- Lee GM, Neutra RR, Hristova L, Yost M, Hiatt RA. 2002. A nested case-control study of residential and personal magnetic field measures and miscarriages. *Epidemiology* 13(1):21-31.
- Lee TM, Ho SM, Tsang LY, Yang SH, Li LS, Chan CC, Yang SY. 2001. Effect on human attention of exposure to the electromagnetic field emitted by mobile phones. *Neuroreport* 12(4):729-31.
- Leszczynski D, Joenväärä S, Reivinen J, Kuokka R. 2002. Non-thermal activation of hsp27/p38MAPK stress pathway by mobile phone radiation in human endothelial cells: Molecular mechanism for cancer- and blood-brain barrier-related effects. *Differentiation* 70:120-129.
- Leszczynski D, Nylund R, Joenväärä S, Reivinen J. 2004. Applicability of discovery science approach to determine biological effects of mobile phone radiation. *Proteomics* 4:426-431.
- Levallois P. 2002. Hypersensitivity of human subjects to environmental electric and magnetic field exposure: a review of the literature. *Environ Health Perspect* 110 Suppl 4:613-8.
- Levallois P, Neutra R, Lee G, Hristova L. 2002. Study of self-reported hypersensitivity to electromagnetic fields in California. *Environ Health Perspect* 110 Suppl 4:619-23.

- Li DK, Odouli R, Wi S, Janevic T, Golditch I, Bracken TD, Senior R, Rankin R, Iriye R. 2002. A population-based prospective cohort study of personal exposure to magnetic fields during pregnancy and the risk of miscarriage. *Epidemiology* 13(1):9-20.
- Lonne-Rahm S, Andersson B, Melin L, Schultzberg M, Arnetz B, Berg M. 2000. Provocation with stress and electricity of patients with "sensitivity to electricity". *J Occup Environ Med* 42(5):512-6.
- Lönn S. 2004. Mobile phone use and risk of intracranial tumors. Stockholm: Karolinska Institutet.
- Lönn S, Ahlbom A, Hall P, Feychting M. 2004. Mobile phone use and the risk of acoustic neuroma. *Epidemiology* 15:653-659.
- Maier R, Greter SE, Maier N. 2004. Effects of pulsed electromagnetic fields on cognitive processes - a pilot study on pulsed field interference with cognitive regeneration. *Acta Neurol Scand* 110(1):46-52.
- Marino A, Nilsen E, Frilot C. 2003. Nonlinear changes in brain electrical activity due to cell phone radiation. *Bioelectromagnetics* 24:339-346.
- Maskarinec G, Cooper J, Swygert L. 1994. Investigation of increased incidence in childhood leukemia near radio towers in Hawaii: preliminary observations. *J Environ Pathol Toxicol Oncol* 13(1):33-7.
- McKenzie DR, Yin Y, Morrell S. 1998. Childhood incidence of acute lymphoblastic leukaemia and exposure to broadcast radiation in Sydney--a second look. *Aust N Z J Public Health* 22(3 Suppl):360-7.
- Michelozzi P, Capon A, Kirchmayer U, Forastiere F, Biggeri A, Barca A, Perucci CA. 2002. Adult and childhood leukemia near a high-power radio station in Rome, Italy. *Am J Epidemiol* 155(12):1096-103.
- Muscat JE, Malkin MG, Shore RE, Thompson S, Neugut AI, Stellman SD, Bruce J. 2002. Handheld cellular telephones and risk of acoustic neuroma. *Neurology* 58(8):1304-6.
- Muscat JE, Malkin MG, Thompson S, Shore RE, Stellman SD, McRee D, Neugut AI, Wynder EL. 2000. Handheld cellular telephone use and risk of brain cancer. *Jama* 284(23):3001-7.
- National Board of Occupational Safety and Health, National Board of Housing Building and Planning, National Electrical Safety Board, National Board of Health and Welfare, Radiation Protection Institute. 1996. Low frequency electrical and magnetic fields: the precautionary principle for national authorities - guidance for decision-makers. Stockholm, Sweden.
- Navarro EA, Segura J, Portolés M, Gómez-Perretta de Mateo C. 2003. The microwave syndrome: a preliminary study in Spain. *Electromagnetic biology and medicine* 22:161-69.
- NRPB. 2003. Health effects from radiofrequency electromagnetic fields. Report of an independent advisory group on non-ionizing radiation.: Vol 14, No. 2: NRPB.
- Nylund R, Leszczynski D. 2004. Proteomics analysis of human endothelial cell line EA.hy926 after exposure to GSM 900 radiation. *Proteomics* 4(5):1359-65.
- Oftedal G, Nyvang A, Moen BE. 1999. Long-term effects on symptoms by reducing electric fields from visual display units. *Scand J Work Environ Health* 25(5):415-21.
- Oftedal G, Wilen J, Sandstrom M, Mild KH. 2000. Symptoms experienced in connection with mobile phone use. *Occup Med (Lond)* 50(4):237-45.
- Peyman A, Gabriel C. Dielectric properties of aging rat tissues at low frequencies.; 2003; Bioelectromagnetics Society 25t Annual meeting, Hawaii, USA. p 378.
- Peyman A, Rezazadeh AA, Gabriel C. 2001. Changes in the dielectric properties of rat tissue as a function of age at microwave frequencies. *Phys Med Biol* 46(6):1617-29.
- Preece AW. unpublished. Investigation of GSM signals on child cognition.

- Repacholi MH, Basten A, Gebiski V, Noonan D, Finnie J, Harris AW. 1997. Lymphomas in E mu-Pim1 transgenic mice exposed to pulsed 900 MHz electromagnetic fields. *Radiat Res* 147(5):631-40.
- Rodier PM. 2004. Environmental causes of central nervous system maldevelopment. *Pediatrics* 113(4 Suppl):1076-83.
- Roosli M, Moser M, Baldinini Y, Meier M, Braun-Fahrlander C. 2004. Symptoms of ill health ascribed to electromagnetic field exposure--a questionnaire survey. *Int J Hyg Environ Health* 207(2):141-50.
- Sandstrom M, Lyskov E, Hornsten R, Hansson Mild K, Wiklund U, Rask P, Klucharev V, Stenberg B, Bjerle P. 2003. Holter ECG monitoring in patients with perceived electrical hypersensitivity. *Int J Psychophysiol* 49(3):227-35.
- Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. 2002a. [Investigation on the health of people living near mobile telephone relay stations: I/Incidence according to distance and sex]. *Pathol Biol (Paris)* 50(6):369-73.
- Santini R, Seigne M, Bonhomme-Faivre L, Bouffet S, Defrasne E, Sage M. 2002b. Symptoms experienced by users of digital cellular phones: a study of a French engineering school. *Electromagnetic biology and medicine* 21:81-88.
- Schuz J, Mann S. 2000. A discussion of potential exposure metrics for use in epidemiological studies on human exposure to radiowaves from mobile phone base stations. *J Expo Anal Environ Epidemiol* 10(6 Pt 1):600-5.
- Sienkiewicz Z, Blackwell R, Haylock R, Saunders R, Cobb B. 2000. Low-level exposure to pulsed 900 MHz microwave radiation does not cause deficits in the performance of a spatial learning task in mice. *Bioelectromagnetics* 21:151-158.
- Swanbeck G, Bleeker T. 1989. Skin problems from visual display units. Provocation of skin symptoms under experimental conditions. *Acta Derm Venereol* 69(1):46-51.
- Swedish Radiation Protection Authority. 2003. Recent Research on Mobile Telephony and Cancer and Other Selected Biological Effects: First annual report from SSI's Independent Expert Group on Electromagnetic Fields. Dnr 00/1854/02, Available at: [http://www.ssi.se/english/EMF\\_exp\\_Eng\\_2003.pdf](http://www.ssi.se/english/EMF_exp_Eng_2003.pdf).
- Thurai M, Goodridge VD, Sheppard RJ, Grant EH. 1984. Variation with age of the dielectric properties of mouse brain cerebrum. *Phys Med Biol* 29(9):1133-6.
- Thurai M, Steel MC, Sheppard RJ, Grant EH. 1985. Dielectric properties of developing rabbit brain at 37 degrees C. *Bioelectromagnetics* 6(3):235-42.
- Wang B, Lai H. 2000. Acute exposure to pulsed 2450-MHz microwaves affects water-maze performance of rats. *Bioelectromagnetics* 21:52-56.
- Ventura C, Maioli M, Asara Y, Santoni D, Mesirca P, Remondini D, Bersani F. 2004. Turning on stem cell cardiogenesis with extremely low frequency magnetic fields. *FASEB* in press.
- Villa S, Ljung L. 1993. Magnetic fields and cancer [Swedish]. Stockholm, Sweden: National Electric and Safety Board.
- Zwamborn A, Vossen S, van Leersum B, Ouwens M, Mäkel W. 2003. Effects of global communication system radio-frequency fields on well being and cognitive functions of human subjects with and without subjective complaints. The Hague, Netherlands: TNO Physics and Electronics Laboratory (TNO-report FEL-03-C148). Available from: URL: [www.ez.nl](http://www.ez.nl). See also [www.gr.nl/pdf.php?ID=1042](http://www.gr.nl/pdf.php?ID=1042). Report nr TNO-report FEL-03-C148.